



A Predominant Urodynamic Diagnosis Can Hide a Minor One: Study in Non-Neurologic Women

F. A. Valentini ^{a*}, B. G. Marti ^a, G. Robain ^a
and P. E. Zimmern ^b

^a Sorbonne Université, Hôpital Rothschild, 75012 Paris, France.

^b UT Southwestern Medical Center in Dallas, TX 75390, USA.

Authors' contributions

This work was carried out in collaboration among all authors. Author FAV did the concept, design of the study, data collection, data analysis and prepare the manuscript. Author BGM collected and analysis the data. Author GR collected the data and prepare the manuscript. Author PEZ concept, design of the study, data analysis and prepare the manuscript. All authors reviewed the results and approved the final version of the manuscript.

Article Information

DOI: 10.9734/JAMMR/2023/v35i185117

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102112>

Original Research Article

Received: 01/05/2023

Accepted: 01/07/2023

Published: 13/07/2023

ABSTRACT

Aims: Voiding dysfunction is a common condition among women. Since voiding and storage symptoms can coexist, evaluation necessitates further investigations with urodynamic studies. Unfortunately, some predominant dysfunction can hide a minor one.

The purpose of our study was to retrospectively review urodynamic records of non-neurologic women referred for evaluation of lower urinary tract dysfunction and to explain hidden concomitant urodynamic diagnoses that might have gone unnoticed without a thorough examination.

Methods: Urodynamic tracings of 404 consecutive non-neurologic women referred for evaluation of lower urinary tract symptoms were reviewed. Initial urodynamic diagnosis had been proposed

*Corresponding author: E-mail: favalentini@gmail.com;

according with ICS/IUGA recommendations and a choice of specific urodynamic criteria. Concomitant urodynamic diagnoses were sought by analyzing the values of characteristic parameters which were hidden by predominant phenomenon.

Results: Concomitant diagnoses were found for 120 (29.7%) women. Coexistent diagnoses were 48 bladder outlet obstruction, 16 detrusor underactivity and 56 intrinsic sphincter deficiency. That condition was observed for women with predominant diagnosis of detrusor overactivity (63.4%) and detrusor overactivity with impaired contractility (60.0%).

Conclusions: Our study show a high prevalence of possible concomitant urodynamic diagnoses for non-neurologic women referred for evaluation of lower urinary tract dysfunction. The practitioner must pay attention to all the parameters measured in order to derive the correct urodynamic diagnosis from which the best management can be proposed.

Keywords: Urodynamic diagnosis; concomitant urodynamic diagnoses; lower urodynamic tract dysfunction; women; non-neurologic.

1. INTRODUCTION AND OBJECTIVES

Lower urinary tract dysfunction (LUTD) is a common condition among women. Non-neurologic voiding dysfunction affects more than 6% of women over 40 [1]. These patients present with a spectrum of different urinary symptoms, related to both storage and voiding, which may be multifactorial in origin or be related to one another. Incontinence (leaking urine) and other urination-related problems belong to a broad category of disorders called voiding dysfunctions. Although common, these conditions can be difficult to discuss and can dramatically diminish quality of life. Voiding dysfunction can be related to detrusor overactivity (DO), detrusor underactivity (DU), bladder outlet obstruction (BOO) and urethral incompetence (ISD). Since voiding and storage symptoms can coexist, they necessitate further evaluation with urodynamics studies (UDS). The goal of UDS is first to evaluate bladder and urethra behavior during the micturition cycle (filling, storage and void), then to propose a diagnosis of LUTD in order to guide towards the best management of the identified dysfunction. Urodynamic criteria for diagnosis are mainly based on measures of pressure (vesical, abdominal and their difference which is the detrusor pressure). Typically, low detrusor pressure with normal flow is associated with unobstructed condition, high pressure with low flow with obstructed condition and low detrusor pressure with low flow with poor detrusor contractility.

But some predominant dysfunction can hide a minor one. Women with detrusor-underactivity-like urodynamic profiles could have concomitant occult bladder outlet obstruction [2]. Resnick first described the coexistence of “detrusor hyperreflexia” and impaired contractility

(DHIC) [3]; that concomitant association is included in the main categories of urodynamic diagnoses according to ICS/IUGA recommendations [4]. Recently COUB a clinical syndrome identified as the coexisting overactive-underactive bladder has been reported which may not be the simple combination of both syndromes [5,6,7]. However, to our knowledge, there has been no study devoted to analysis of hidden concomitant urodynamic diagnoses.

The purpose of our study was to retrospectively review urodynamic records of non-neurogenic women referred for evaluation of LUTD and to explain how hidden concomitant urodynamic diagnosis (UD) that might have gone unnoticed without a thorough examination had been detected.

2. METHODS

Urodynamic records of consecutive women referred for evaluation of LUTD were analyzed. Criteria for inclusion were PFs tracings providing maximum flow rate Q_{max} and detrusor pressure at Q_{max} ($p_{det.Q_{max}}$) without significant contribution of abdominal pressure between onset of flow and Q_{max} ($< 3 \text{ cm H}_2\text{O}$), a $Q_{max} \geq 2 \text{ mL/s}$, a voided volume $\geq 100 \text{ mL}$, and a non-interrupted flow.

Criteria for exclusion were neurological disease, diabetes mellitus, stage ≥ 2 prolapse and failure to understand simple orders or a Mini-Mental State score < 20 .

All patients were evaluated using medical history, review of medications, bladder diary for at least 3 days including voiding times and voided volumes both day and night, physical examination, and dipstick urinalysis to eliminate urinary tract infection. Urodynamic sessions were performed

by the same team using the Dorado® unit from Laborie. Urodynamic tests were carried out according to the International Continence Society Good Urodynamic Practices [8]. Cystometry was performed with the patient in the sitting position. Bladder was filled with saline at room-temperature at a medium filling rate of 50 mL/min. Filling cystometrogram was obtained via a triple lumen urethral catheter 7 F allowing for urethral pressure recording, followed by an intubated flow (IF). Pressure transducers were zeroed to atmospheric pressure at the upper edge of the symphysis pubis. Rectal pressure was recorded using a punctured intrarectal balloon catheter filled with 2 mL of saline according to the report of Good Urodynamic Practice guidelines [9]. Urethral pressure profilometry (UPP) was performed after IF, at rest, bladder filled at 200 mL; Valsalva maneuver was added to search for intrinsic sphincter deficiency (ISD).

According to ICS/IUGA recommendations [4] the main categories of urodynamic diagnoses were bladder outlet obstruction (BOO), detrusor hyperactivity with impaired contractility (DHIC), detrusor overactivity (DO), or detrusor underactivity (DU). Some investigations were found “normal” (N) and other related to urethral dysfunction: ISD or voiding triggered by urethral relaxation [URA].

Each predominant urodynamic diagnosis was associated with specific urodynamic parameters values that were routinely used in our department.

- For BOO, cut-off values proposed by Defreitas and colleagues [10] were chosen: $Q_{max} < 12$ mL/s and $p_{det.Qmax} > 25$ cm H₂O. Comparison was made with the value of Solomon-Greenwell index $BOOI = (p_{det.Qmax} - 2.2 * Q_{max})$ [11]: likely obstructed when BOOI was higher than 5 and obstruction almost certain when BOOI was higher than 18.
- DHIC mainly observed in elderly patients implied detrusor overactivity during storage and impaired emptying with low flow of long duration leading to possible high PVR [3].
- DO was the observation of detrusor contractions of varying durations and amplitudes during filling cystometry. DO was classified as phasic (during filling cystometry) and terminal (a single involuntary detrusor contraction that

cannot be suppressed and that occurs when the maximum bladder capacity is reached, resulting in urinary incontinence and often complete bladder emptying) [12].

- For DU, which is characterized by low pressure-low flow, cutoff values proposed by Gammie et al. [13] were chosen: $p_{det}@Qmax < 20$ cmH₂O, $Q_{max} < 15$ mL/s and $BVE\% < 90$.
- N was associated with no abnormality detected during PFs.
- ISD was defined as low MUCP (maximum urethral closure pressure) vs age ((120 – age) - 20%) [14] or MUCP less than 20/35 cm H₂O [15] and/or positive VLPP less than 60 cm H₂O [16].
- URA was defined as voiding triggered by urethral relaxation (both urethral and flow curves recordings).

When detrusor contractility was needed to verify one proposal, VBN detrusor contractility parameter k [17] (without unit) and PIP1 index ($PIP1 = p_{det.Qmax} + Q_{max}$) [18] (in cm H₂O) were calculated.

Minor UD were suspected when specific urodynamic parameters values were found close to the values allowing predominant UD were observed.

This retrospective study was conducted in accordance with the declaration of Helsinki. The local practice of our Ethics Committee does not require a formal institutional review board approval for retrospective studies.

2.1 Statistical Analysis

Data are presented as mean ± SD and range. Analysis of variance (ANOVA), t-test, and the chi-square test were used as appropriate. All statistical results were considered significant at $p < 0.05$. Statistical analyses were performed using SAS, version 5.0 (SAS Institute, Inc., Cary, NC).

3. RESULTS

Over a period of 8 years, 404 urodynamic tracings of consecutive non-neurogenic women referred for LUTS met study criteria. The urodynamic study was performed by the same team over time. Mean age was 58.4 ± 16.5 years [20-96 y]. Main complaint was urinary incontinence: 87 stress (SUI), 122 mixed (MUI), and 111 urge (UI). Forty four women with

complaints other than incontinence were called OTHER; among them, 33 complained of urinary frequency (PK) and 14 from dysuria (Dys) (Table1).

Among predominant diagnoses, some coexistent diagnoses were found (Table 2).

That co-existent condition was found in 120 (29.7%) women and was mainly observed in women with predominant diagnosis of DO (63.4%) and DHIC (60.0%).

Concomitant UD of BOO according to Defreitas' criteria [9] was found in 48 women first diagnosed as DHIC (3), DO (37) and ISD(8); among them, 20 (41.6%) had obstruction almost certain, 24 (50.0%) were likely obstructed and 4(8.3%) non-obstructed according to the Solomon-Greenwell index [10]. Values of contractility parameters were higher in patients with only BOO diagnosis ($k = .81 \pm .58$; $PIP1 = 55.8 \pm 22.5$) vs those with concomitant BOO and ISD ($k = .63 \pm .37$; $PIP1 = 48.7 \pm 18.4$).

Concomitant UD of DU according with Gammie's criteria [13] was found in 16 women with the first diagnosis of DHIC (2), DO (3), ISD (8) and URA (3).

Concomitant UD of ISD was found in 56 women first diagnosed as BOO (10), DHIC (7), DO (14), DU (22) and URA (3).

For some women, two coexistent diagnoses were observed: 11 women (predominant UD: 2 DHIC and 9 DO) had two concomitant UD which were BOO and ISD and one woman (predominant UD was DO) had two concomitant UD which were ISD and DU.

4. DISCUSSION

LUTD express as LUTS which are considered to represent a multifactorial constellation of non-specific symptoms that affect bladder function during both filling and voiding.

Symptoms such as stress urinary incontinence (SUI) and detrusor overactivity (DO) can co-exist in a number of patients. A non-infrequent situation is the co-existence of irritative (urgency, nocturia ...) and obstructive (incomplete voiding, low flow...) LUTS. So, high PVR can be due to

detrusor underactivity (DU), bladder outlet obstruction (BOO) or a combination of both.

Recently, some studies have been devoted to the study of coexistent syndromes (COUB) [5-6-7] but to our knowledge there is no study which analyzes the possible concomitant urodynamic diagnosis in a non-neurogenic female population and the authors say that "Invasive urodynamic tests may be necessary in unclear cases or in cases not responding to initial treatment of the most troublesome symptoms".

Detailed assessment of bladder function using more complex urodynamic studies is needed in any patient where previous therapy has proved unsuccessful. But different urodynamic findings may be present within a given clinical presentation, and the same urodynamic observations may be made in the presence of different symptoms. Moreover, due to the physiological variability of lower urinary tract function, there are some limits to urodynamic investigation.

But we can observe that some of these diagnoses are concomitant with some other possible diagnoses when reviewing other UDS parameters values. Thus secondary urodynamic diagnoses are based on observation of values of characteristic parameters which are hidden by predominant phenomenon such as uninhibited detrusor contractions during filling (DHIC, DO) or combination of parameters values retained as characteristic of diagnosis, like DU. It is the case when there are low MUCP or/and VLPP < 60cm H₂O characteristic of ISD with predominant diagnosis of DHIC, DO and DU.

An interesting observation is that we can compare Defreitas' criteria and Solomon-Greenwell index. Among patients with predominant diagnosis BOO according to Defreitas'criteria 72.7% were diagnosed obstruction almost certain according to Solomon-Greenwell index. Furthermore, when obstruction was almost certain according to Solomon-Greenwell index, it was found in 72.7% in predominant diagnosis and only 34.5% when BOO is proposed as secondary diagnosis using the Defreitas criteria. So, these criteria/indices for BOO appear valuable despite the fact that there is no widely accepted precise definition for that condition without confirmation of the obstruction site by imaging (voiding cystogram or videourodynamic study).

Table 1. Main complaint vs. predominant urodynamic diagnosis

	BOO	DHIC	DO	DU	ISD	N	URA	Nbr
SUI	10	2	8	13	26	19	9	87
MUI	9	8	27	17	25	32	4	122
UUI	14	7	37	13	11	27	2	111
OTHER	11	3	10	17	7	29	7	84
Nbr	44	20	82	60	71	105	22	404

*SUI: Stress Urinary Incontinence; MUI: Mixed Urinary Incontinence; UUI: Urge Urinary Incontinence; OTHER: Urinary Complaint without Urinary Incontinence.
BOO: Bladder Outlet Obstruction; DHIC: Detrusor Overactivity with Impaired Contractility; DO Detrusor overactivity; DU: detrusor underactivity; ISD: intrinsic sphincter deficiency; N investigations found "normal":
URA: voiding triggered by urethral relaxation.
Nbr: Number of Patients*

Table 2. Concomitant diagnosis vs. predominant urodynamic diagnosis (Nbr-%)

	BOO	DHIC	DO	DU	ISD	N	URA
Nbr-	44	20	82	60	68	105	22
%	(10.9%)	(4.9%)	(20.3%)	(14.8%)	(16.8%)	(26.0%)	(5.4%)
BOO/BOOIf>18	0	1(5.0%)	19(23.2%)	0	0	0	0
5<BOO/BOOIf<18	0	2(10.0%)	16(19.5%)	0	6(8.8%)	0	0
DU	0	2(10.0%)	3(3.6%)	0	8(11.7%)	0	3(13.6%)
ISD	10(22.7%)	7(35%)	14(17.1%)	22(36.7%)	0	0	3(13.6%)
Σ % concomitant diagnosis	22.7%	60.0%	63.4%	36.7%	20.6%	0	27.3%

*BOOIf: Bladder Outlet Obstruction Index female
BOO: Bladder Outlet Obstruction; DHIC: Detrusor Overactivity With Impaired Contractility; DO: Detrusor Overactivity; DU: Detrusor Underactivity; ISD: Intrinsic Sphincter Deficiency; N investigations found "normal":
URA: Voiding Triggered Byurethral Relaxation.
Nbr: Number of Patients*

Secondary UD of ISD is the most frequently observed in women with predominant UD of DHIC and DU. That observation is mainly due to aging. Women with these predominant UD are older, respectively 65 and 67 years which explain the low MUCP vs age observed.

ISD is present when predominant UD is BOO while the concerned population is younger (mean age 55 years old) and without neurologic pathology. That result could be an unexpected consequence of obstruction leading to an impaired sphincter function. That hypothesis is supported by the decrease of the values in contractility parameters.

A similar explanation can be proposed for occurrence of secondary UD of ISD for DO patients while important muscular relaxation is most likely explanation for URA patients.

Thus, urodynamic diagnosis must be proposed with circumspection in intricaded clinical presentations and need the strict evaluation of all information obtained from pre urodynamic testing evaluation.

First limitation of this study is that it is retrospective and from a single-center. Second limitation is the choice of specific urodynamic criteria to give each predominant urodynamic diagnosis although for some there is no consensus. Third limitation is that the studied population includes women referred to our urodynamic laboratory for evaluation of LUTS; the physician performing the urodynamic investigation was not the physician who originally requested urodynamics. Therefore, our findings can only be considered as advice. Lastly, there are limitations to the use of k as a detrusor contractility index, and those are primarily related to the voiding performance. As already alluded to, the limitations for k interpretation include a non-interrupted flow, a voided volume > 100 mL, $Q_{max} > 2\text{mL/s}$ and no significant abdominal straining (to our knowledge these conditions have not been evaluated in the development of some other indices). However, the software allowing evaluation of k (in Excel) is easy to use and can be obtained (with instructions) on request from its authors.

5. CONCLUSION

Our study show a high prevalence of possible concomitant urodynamic diagnoses for non-neurologic women referred for evaluation of lower urinary tract dysfunction. Although the criteria generally accepted to render a urodynamic diagnosis are useful, the practitioner must pay attention to all the parameters measured and pay close attention to diagnoses that might go unnoticed without a thorough examination in order to derive the correct urodynamic diagnosis from which the best management can be proposed.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Raheem AA, Madersbacher H. Voiding dysfunction in women: How to manage it correctly. *Arab J Urol.* 2013;11:319–330. DOI: 10.1016/j.aju.2013.07.005
2. Chow PM, Hsiao SM, Kuo HC. Identifying occult bladder outlet obstruction in women with detrusor-underactivity-like urodynamic profiles. *Sci Rep.* 2021;11:232-42. DOI: 10.1038/s41598-021-02617-0
3. Resnick NM, Yalla SV. Detrusor or hyperactivity with impaired contractile function. An unrecognized but common cause of incontinence in elderly patients. *JAMA.* 1987;257:3076-81. DOI: 10.1001/jama.257.22.3076
4. Haylen BT, de Ridder D, Freeman RM, Swift SE, Berghmans B, Lee J, Monga A, Petri E, Rizk DE, Sand PK, Schaer GN. An international urogynecological association (IUGA)/international continence society (ICS) joint report on the terminology for female pelvic floor dysfunction. *Neurourol Urodyn.* 2010;29:4-20. DOI: 10.1002/nau.20798
5. Gammie A, Kaper M, Steup A, Yoshida S, Dorrepaal C, Kos T, Abrams P. What are the additional signs and symptoms inpatients with detrusor underactivity and coexisting detrusor overactivity? *Neurourol Urodyn.* 2018;37:2220–2225. DOI: 10.1002/nau.23565
6. Mancini V, Tarcan T, Serati M, Wyndaele M, Carrieri G, Abrams P. Is coexistent overactive–underactive bladder (with or without detrusor overactivity and underactivity) a real clinical syndrome? *ICI-RS 2019.Neurourol Urodyn.* 2020; 39: S50-S59. DOI: 10.1002/nau.24311
7. Frigerio M, Barba M, Marino G, Volonte S, Melocchi T, De Vicari D, Torella M, Salvatore S, Braga A, Serati M, Manodoro S, Cola A. Coexistent Detrusor Overactivity- Underactivity in Patients with Pelvic Floor Disorders. *Healthcare.* 2022; 10:1720. DOI: 10.3390/healthcare10091720
8. Rosier PFWM, Schaefer W, Lose G, Goldman HB, Guralnick M, Eustice S, Dickinson T, Hashim H. International continence society good urodynamic practices and terms 2016: urodynamics, uroflowmetry, cystometry, and pressure-flow study. *Neurourol Urodyn.* 2017; 36: 1243-60, dx.doi.org/10.1002/nau.23124.
9. Schäfer W, Abrams P, Liao L, Mattiasson A, Pesce F, Spanberg A, Sterling AM, Zinner NR, van Kerrebroeck P. Good urodynamic practices: uroflowmetry, fillingcystometry, and pressure-flow studies. *Neurourol Urodyn.* 2002; 21:261-74.
10. Defreitas GA, Zimmern PE, Lemack GE, Shariat SF. Refining diagnosis of anatomic female bladder outlet obstruction: comparison of pressure-flow study parameters in clinically obstructed women with those of normal controls. *Urology.* 2004; 64: 675-9 doi: 10.1016/j.urology.2004.04.089.
11. Solomon E, H, Duffy M, Rashid T, Akinluyi E, Greenwell TJ. Developing and validating a new nomogram for diagnosing bladder outlet obstruction in women. *Neurourol Urodyn.* 2017; 37 : 368-78 DOI: 10.1002/nau.23307
12. Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, van Kerrekroek P, Victor A, Wein A. The standardisation of terminology of lower urinary tract function: Report from the Standardisation Sub-

- committee of the International Continence Society. *Neurourol Urodyn.* 2002;21:167-178.
DOI: 10.1002/nau.10052
13. Gammie A, Kaper M, Dorrepaal C, Kos T, Abrams P. Signs and symptoms of detrusor underactivity: An analysis of clinical presentation and urodynamic tests from a large group of patients undergoing pressure flow studies. *Eur Urol.* 2016;69: 361–369.
DOI: 10.1016/j.euro.2015.08.014
 14. Rud T. Urethral pressure profile in continent women from childhood to old age. *Acta Obstet Gynecol Scand.* 1980;59: 331-335.
 15. Kapoor DS, Housami F, White P, Swithinbank L, Drake M. Maximum urethral closure pressure in women: Normative data and evaluation as a diagnostic test. *Int UrogynecolJ.* 2012;23:1613–1618.
DOI: 10.1007/S00192-012-1770-7
 16. Burden H, Warren K, Abrams P. Leak point pressures: How useful are they? *Curr Opin Urol.* 2005;25:317-322
 17. Valentini FA, Besson GR, Nelson PP, Zimmern PE. A mathematical micturition model to restore simple flow recordings in healthy and symptomatic individuals and enhance uroflow interpretation. *Neurourol Urodyn.* 2000;19(2):153-176.
 18. Tan TL, Bergmann MA, Griffiths D, Resnick NM. Stop test or pressure-flow study? Measuring detrusor contractility in older females. *Neurourol Urodyn.* 2004;23: 184-9.
DOI: 10.1002/nau200.20

© 2023 Valentini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102112>