



## Evaluation of *In vitro* Nematicidal Efficiency of Copper Nanoparticles against Root-knot Nematode *Meloidogyne Incognita*

Elwy A. Mohamed<sup>1\*</sup>, Sherif F. Elsharabasy<sup>1</sup> and Doaa Abdulsamad<sup>2</sup>

<sup>1</sup>Central Laboratory for Date Palm Research and Development, Agricultural Research Center, P.O.Box 12619, Giza, Egypt.

<sup>2</sup>Department of Biotechnology, October University for Modern Sciences and Technology, P.O.Box 12585, Giza, Egypt.

### Authors' contributions

This work was carried out in collaboration between all authors. Author EAM designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors SFE and DA managed the literature searches and the analyses of the study. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/SAJP/2019/46065

#### Editor(s):

(1) Dr. Ashlesha Kaushik, Department of Pediatric Infectious Diseases, Unity Point Health, United States.

#### Reviewers:

(1) Mahmoud M. A. Youssef, National Research Centre, Egypt.

(2) Muhammad Shahzad Aslam, Xiamen University, Malaysia.

(3) ABA-Toumou Lucie, University of Bangui, Central African Republic.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46065>

Original Research Article

Received 19 October 2018

Accepted 03 January 2019

Published 14 January 2019

### ABSTRACT

Root-knot nematodes (*Meloidogyne* spp.) are considered among the most deteriorating soilborne parasites that can significantly affect many plants. These nematodes are developing increased resistance against nematicides used currently to control them, therefore, continued use of these nematicides poses a challenge, thereby giving rise to the need for newer alternatives. This paper evaluated the *In vitro* nematicidal efficiency of copper nanoparticles (CuNPs) against root-knot nematode, *Meloidogyne incognita*. In this study, CuNPs were prepared according to the chemical reduction method; physicochemical characterization of CuNPs was done using UV-Vis spectroscopy, Dynamic Light Scattering and Transmission Electron Microscopy. When second stage juveniles (J2) of *M. incognita* were incubated in soil saturated with CuNPs (spherical shape; 100 nm diameter) for 3 days, it was found that J2 mortality is directly proportional to the

\*Corresponding author: E-mail: nano.elwy@gmail.com;

concentration of CuNPs and 0.2 g/L was sufficient to cause 100% mortality. Statistical analysis showed that all mortalities caused by treatment with CuNPs at different concentrations were statistically significant compared with non-treated control. Conclusively, this paper may provide a potential alternative nematicide against root-knot nematode *Meloidogyne incognita*. Further *In vivo* and toxicological research on CuNPs should be conducted in order to assess the possible applicability of such nanoparticles as a nematicide.

**Keywords:** *Meloidogyne incognita*; nematicide; copper nanoparticles; nematicide alternative.

## 1. INTRODUCTION

*Meloidogyne* spp. was first reported in cassava (*Manihot esculenta*) by Neal in 1889 [1]; Since then, root-knot nematodes (*Meloidogyne* spp.) are considered among the most deteriorating soilborne parasites that can significantly affect many field crops, trees and turfgrass [2]. Nematodes are characterized with a broad host range of greater than 3,000 plant species [3]. Furthermore, it was reported that around 5% of the world crop production was lost annually due to infection with *Meloidogyne* species [4] and the losses can reach up to 64% of the yield [5-7].

Negative effects of nematode infections are not limited to decreased productivity of the economical crops, since it can also affect the playability and aesthetic quality of golf courses [8].

*Meloidogyne* species encompass 98 species, among them *M. incognita*, *M. javanica*, *M. hapla*, and *M. arenaria* are considered the most common [9].

After banning Nematicur in 2008 due to environmental concerns, there is a dire need for developing newer efficient alternatives to control such plant-parasitic nematodes. In this respect, the narrow range effectiveness characterizing biological control agents limits its applicability. For example, the bacterial parasite, *Pasteuria* sp. can control sting nematodes (*Belonolaimus longicaudatus*) [10]; however, it cannot affect the other species of plant-parasitic nematodes such as root-knot nematodes (*Meloidogyne* spp.).

Nanotechnology is considered a promising and effective means for controlling root-knot nematode, wherein some papers reported the nematicidal effect of silver nanoparticles (AgNPs) [11,12], gold nanoparticles [13] and silicon carbide nanoparticles [14] against root-knot nematodes. By virtue of the well-established nematicidal effect of AgNPs, AgNPs were proposed [15] as a potential alternative nematicide.

In this regard, many papers have established a robust emphasis on the antimicrobial effect of CuNPs [16-18]; thus, in this paper, we evaluate the *In vitro* nematicidal efficiency of CuNPs against J2 *M. incognita* as another potential alternative for controlling such parasites.

## 2. MATERIALS AND METHODS

### 2.1 Chemicals

All chemicals used were analytical grade of purity and were used without further purification.

L-ascorbic acid (Future Modern Co., Egypt.); Cetyl trimethylammonium bromide (CTAB) (Sigma-Aldrich, Egypt.); copper sulfate pentahydrate (Elnasr Pharmaceuticals Co., Egypt) were used to prepare CuNPs.

### 2.2 Methods

#### 2.2.1 Preparation of CuNPs

CuNPs were prepared according to the chemical reduction method [19]. In this method, L-ascorbic acid was used as a reducing agent, in the presence of CTAB as a cationic surfactant, to reduce copper cations provided from copper sulfate pentahydrate into copper atoms, which were aggregated and developed into copper nanoparticles, with their characteristic reddish brown color, at pH of 6.8 and temperature of 85°C. CuNPs were centrifugally (4000 rpm) collected for further characterization and application.

#### 2.2.2 Characterization of CuNPs

The characteristic surface plasmon resonance of the synthesized CuNPs was detected using UV-Vis Spectrophotometer (ORION AQUAMATE 8000). Also, particles size distribution by number of CuNPs was detected using Dynamic light scattering (DLS) (Zetasizer nano series (Nano ZS), Malvern, UK). Moreover, the shape of the CuNPs was detected through Transmission Electron Microscopy (Tecnai G20, Super twin, double tilt, FEI, Netherland).

### 2.2.3 *In vitro* application of CuNPs

21 jars (300 cm<sup>3</sup>) were filled with soil composed of 1:1 beet moss and sand. Water saturation level of 300 cm<sup>3</sup> soil was determined to be 100 ml. Each filled jar was inoculated with 1,000 larva second stage juveniles (J2) and homogenized well. Then, each jar was saturated with 100 ml of copper nanoparticles solution at different concentrations, (0.02, 0.04, 0.06, 0.08, 0.1 and 0.2 g/L). Soil jars saturated with water were used as a control. Each concentration was applied in triplicate. All jars were incubated at room temperature for 3 days. After the mentioned exposure time, nematodes were extracted, counted and mortality was calculated according to equation (1).

$$\text{Mortality (\%)} = \left( \frac{\text{Number of Dead Nematodes}}{\text{Total Number of Nematodes}} \right) \times 100 \quad (1)$$

### 2.3 Statistical Analysis

SPSS 22 software (Chi Square Method) was used at  $P \leq 0.05$  to distinguish between the nematicidal efficacies. Each treatment was conducted in triplicate, and the whole experiment was repeated twice [20].

## 3. RESULTS

### 3.1 Physicochemical Characterization of CuNPs

Successful synthesis of CuNPs was confirmed through exhibiting their characteristic surface plasmon resonance peak which was detected using UV-Vis Spectrophotometer (ORION

AQUAMATE 8000) at wavelength of 572 nm [21], as shown in Fig. 1.

Also, Dynamic Light Scattering revealed that the average size of the synthesized CuNPs was about 100 nm; as shown in Fig. 2.

In addition, Transmission Electron Microscopy revealed that the synthesized CuNPs have spherical shape, as shown in Fig. 3.

### 3.2 Evaluation of the Nematicidal Effect of CuNPs

Statistical analysis showed that all concentrations of CuNPs exhibited significant inhibitions of the J2 *M. incognita*. It was shown that CuNPs have a linear nematicidal effect against J2 *M. incognita*, i.e. the higher the concentration of CuNPs, the higher the mortality of nematodes. The concentration of 0.2 g/L was sufficient to completely inactivate all nematodes. Viable nematodes are circular or curved, while dead nematodes are straight, as shown in Fig. 4.

Concentration-dependent mortality of *M. incognita* caused by CuNPs can be shown in Fig. 5.

## 4. DISCUSSION

This study has emphasized on the potential *In vitro* nematicidal effect exhibited by CuNPs against the second stage juveniles (J2) of root-knot nematodes, *M. incognita*; this was demonstrated through the significant increase of J2 mortality at various concentrations of CuNPs compared with non-treated control.






Fig. 1. Characteristic surface plasmon resonance peak of CuNPs at 572 nm



**Fig. 2. Particle size distribution by number of CuNPs, showing the average particle size of about 100 nm**



**Fig. 3. Transmission Electron Micrograph of the synthesized CuNPs showing the spherical shape of the particles**



**Fig. 4. the shape of viable vs. dead nematodes under compound microscope**

Recently, the effect of silicon carbide nanoparticles on hatching and survival of *M. incogneta* was investigated [14]. In that study, it was found that silicon carbide nanoparticles neither affect hatchability of larvae nor survival of second stage juveniles (J2) of *M. incogneta*. Which urge the need for assessing the

nematicidal effect of more toxic nanoparticles against such tolerant nematodes. In this regard, CuNPs may offer that alternative due to their potential nematicidal effect against *M. incogneta*, as shown from the present study.

Also, among different types of nanoparticles, the nematicidal effect of AgNPs has extensively studies; but, from this investigation, it is noteworthy that CuNPs could exhibit a significantly higher nematicidal effect than AgNPs at the same concentration against J2 of root-knot nematodes, *M. incogneta*. In this regard, it was reported that 200 ppm of AgNPs could cause a mortality of 52% at the third day of direct exposure in water [22]. On the other hand, CuNPs at the same concentration could achieve a mortality of 100% after 3 days of indirect exposure in soil. This may due to the profound toxic effect of copper nanoparticles causing DNA damage, which is in contrasts to the more mild effect of AgNPs, which depends mainly on disturbance of many cellular mechanisms such as synthesis of ATP, permeability of the cellular membrane and response to the oxidative stresses in prokaryotes [23,24] and eukaryotes [12,25].

In addition, it was reported [26] that the highest percentage of mortality achieved after 3 days of direct exposure of second stage juveniles (J2) to AgNPs was 95%; while higher mortality percentage (100%) was attained using CuNPs, despite the indirect exposure, which reflect the superior nematicidal efficiency of CuNPs over both silicon carbide nanoparticles and AgNPs. .

Furthermore, the non-specific nematicidal effect of copper nanoparticles provided a relative advantage over the microbial agents of bio-control, which are limited with their relatively high specific host range among different nematode species.

But some concerns may arise due to the emphasized toxicity of CuNPs [27]. In this regard, our paper just confirms the nematicidal effect of copper nanoparticles, this effect can be exploited to control nematodes infecting, for example, ornamental plants in pots or turfgrass, but not to control nematodes infecting, for example, edible crops; so as not to harm the human or environment. Otherwise, further research should be conducted to minimize such toxic effect of CuNPs through, for example, masking CuNPs or loading them on non-toxic matrix such that increase its specific targeting to only nematodes.




Fig. 5. A graph shows the direct proportionality between CuNPs concentration and J2 mortality

## 5. CONCLUSION

To sum up all, it can be concluded that CuNPs may provide an alternative nematicide against the root-knot nematodes, *M. incognita*. But, further research should be conducted in order to investigate the environmental consequences of CuNPs, hence determining the optimum doses and methods that can be applied in field without considerable hazards.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Neal JC. The root-knot disease of the peach, orange and other plants in Florida, due to the work of *Anguillula*. Bulletin US Division of Entomology. 1889;20.
2. Gill HK, Mcsorley R. Cover crops for managing root-knot nematodes. University of Florida, IFAS Extension, ENY-063. 2011;1-6.
3. Reddy DDR. Analysis of crop losses in tomato due to *Meloidogyne incognita*. Indian Journal of Nematology. 1985;15(1): 55-59.
4. Karajeh M. Interaction of root-knot nematode (*Meloidogyne javanica*) and tomato as affected by hydrogen peroxide. Journal of Plant Protection Research. 2008;48(2):181-187.
5. Roberts PA, Matthews WC, Ehlers JD. Root-knot nematode resistant cowpea cover crops in tomato production systems. Agronomy Journal. 2005;97(6): 1626-1635.
6. Sikora RA, Schäfer K, Dababat AA. Modes of action associated with microbially induced in planta suppression of plant-parasitic nematodes. Australasian Plant Pathology. 2007;36(2):124-134.
7. Balbaa EME. Biological and environmental studies on root-knot nematode affecting Leguminosae [M.Sc. Thesis]. Egypt: Faculty of Agriculture, Alexandria University. 2010;50.
8. Crow WT. Plant-parasitic nematodes on golf course turf. Outlooks on Pest Management. 2005;16(1):10-15.
9. Jones JT, Haegeman A, Danchin EG, Gaur HS, Helder J, Jones MG, Kikuchi T, Manzanilla-López R, Palomares-Rius JE, Wesemael WM, Perry RN. Top 10 plant-parasitic nematodes in molecular plant pathology. Molecular Plant Pathology. 2013;14(9):946-961.
10. Luc JE, Pang W, Crow WT, Giblin-Davis RM. Effects of formulation and host nematode density on the ability of *in vitro*-produced *Pasteuria endospores* to control its host *Belonolaimus longicaudatus*. Journal of Nematology. 2010;42(2):87.
11. Roh JY, Sim SJ, Yi J, Park K, Chung KH, Ryu DY, Choi J. Ecotoxicity of silver nanoparticles on the soil nematode, *Caenorhabditis elegans* using functional ecotoxicogenomics. Environmental Science & Technology. 2009;43(10):3933-3940.
12. Lim D, Roh JY, Eom HJ, Choi JY, Hyun J, Choi J. Oxidative stress-related PMK-1 P38 MAPK activation as a mechanism for

- toxicity of silver nanoparticles to reproduction in the nematode, *Caenorhabditis elegans*. Environmental Toxicology and Chemistry. 2012;31(3): 585-592.
13. Thakur RK, Shirkot P, Dhirta B. Studies on effect of gold nanoparticles on *Meloidogyne incognita* and tomato plants growth and development. Bio Rxiv. 2018; 428144.
  14. Al Banna L, Salem N, Ghrair A, Habash S. Impact of silicon carbide nanoparticles on hatching and survival of soil nematodes *Caenorhabditis elegans* and *Meloidogyne incognita*. Applied Ecology and Environmental Research. 2018;16(3):2651-2662.
  15. Cromwell WA, Yang J, Starr JL, Jo YK. Nematicidal effects of silver nanoparticles on root-knot nematode in bermudagrass. Journal of Nematology. 2014;46(3):261.
  16. Karthik AD, Geetha K. Synthesis of copper precursor, copper and its oxide nanoparticles by green chemical reduction method and its antimicrobial activity. Journal of Applied Pharmaceutical Science. 2013;3(5):16.
  17. Betancourt-Galindo R, Reyes-Rodriguez PY, Puente-Urbina BA, Avila-Orta CA, Rodríguez-Fernández OS, Cadenas-Pliego G, Lira-Saldivar RH, García-Cerda LA. Synthesis of copper nanoparticles by thermal decomposition and their antimicrobial properties. Journal of Nanomaterials. 2014;1-5.
  18. Viet PV, Nguyen HT, Cao TM, Hieu LV. *Fusarium antifungal* activities of copper nanoparticles synthesized by a chemical reduction method. Journal of Nanomaterials. 2016;1-7.
  19. Biçer M, Şişman İ. Controlled synthesis of copper nano/microstructures using ascorbic acid in aqueous CTAB solution. Powder Technology. 2010;198(2):279-284.
  20. McDonald JH. Handbook of biological statistics. Sparky House Publishing, Baltimore; 2008.
  21. Kaminskienė Ž, Prosyčėvas I, Stonkutė J, Guobienė A. Evaluation of optical properties of Ag, Cu, and Co nanoparticles synthesized in organic medium. Acta Physica Polonica A. 2013;123(1).
  22. Taha Entsar H. Nematicidal effects of silver nanoparticles on root-knot nematodes (*Meloidogyne incognita*) in laboratory and greenhouse. Journal of Plant Protection and Pathology, Mansoura Univ. 2016;7(5):333–337.
  23. Lok CN, Ho CM, Chen R, He QY, Yu WY, Sun H, Tam PK, Chiu JF, Che CM. Proteomic analysis of the mode of antibacterial action of silver nanoparticles. Journal of Proteome Research. 2006;5(4): 916–924.
  24. Choi O, Hu Z. Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria. Environmental Science & Technology. 2008;42(12):4583-4588.
  25. Ahamed M, Posgai R, Gorey TJ, Nielsen M, Hussain SM, Rowe JJ. Silver nanoparticles induced heat shock protein 70, oxidative stress and apoptosis in *Drosophila melanogaster*. Toxicology and Applied Pharmacology. 2010;242(3):263-269.
  26. Hassan ME, Zawam HS, El-Nahas SE, Desoukey AF. Comparison study between silver nanoparticles and two nematicides against *Meloidogyne incognita* on tomato seedlings. Plant Pathology Journal. 2016; 15(4):144-151.
  27. Karlsson HL, Cronholm P, Gustafsson J, Moller L. Copper oxide nanoparticles are highly toxic: A comparison between metal oxide nanoparticles and carbon nanotubes. Chemical Research in Toxicology. 2008; 21(9):1726-1732.

© 2019 Mohamed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
 The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/46065>