



Micronutrient Dynamics in Soil under Integrated Thermochemical Organic Fertilizer Pellet Application

Phurpa Tsering^{a++*}, Naveen Leno^{a#}, Rani B^{a†},
Sarada S^{b‡} and Raji Swaroop^{a#}

^a Department of Soil Science and Agricultural Chemistry, College of Agriculture, Vellayani, Trivandrum, Kerala, India.

^b Department of Vegetable Science, College of Agriculture, Vellayani, Trivandrum, Kerala, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84892>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121139>

Original Research Article

Received: 30/05/2024
Accepted: 01/08/2024
Published: 07/08/2024

ABSTRACT

A soil incubation study was carried out in the Department of Soil Science and Agricultural Chemistry, to assess dynamics of micronutrient release. Soil pots were filled with 5 kg air-dried, 2 mm sieved soil. The study was laid in Completely Randomized Design with 5 treatments and 4 replications. Treatments were, T₁: Pellet formulation 1; T₂: Pellet formulation 2; T₃: Pellet formulation 3; T₄: FYM+NPK (RDF as per POP) and T₅: Control (Soil alone). The Fe, Mn, Zn, Cu

⁺⁺ PG Scholar;

[#] Assistant Professor;

[†] Professor and Head;

[‡] Assistant Professor and Head;

*Corresponding author: E-mail: phurpa-2021-11-121@student.kau.in;

Cite as: Tsering, Phurpa, Naveen Leno, Rani B, Sarada S, and Raji Swaroop. 2024. "Micronutrient Dynamics in Soil under Integrated Thermochemical Organic Fertilizer Pellet Application". *International Journal of Plant & Soil Science* 36 (8):624-31. <https://doi.org/10.9734/ijpss/2024/v36i84892>.

and B content of soil were 10.14 mg kg⁻¹, 8.59 mg kg⁻¹, 1.90 mg kg⁻¹, 2.50 mg kg⁻¹ and 0.30 mg kg⁻¹, respectively. Three distinct pellet formulations P₁, P₂ & P₃ were used in study based on thermochemical organic fertilizer (TOF). Fe, Mn and Zn exhibited a gradual and progressive release pattern throughout the incubation period. Whereas Cu and B availability showed a decreasing trend from the 15th day onwards. Highest Fe, Mn and Zn availability were observed towards the end of the incubation study. While the Cu and B recorded highest on the initial 15th day of incubation. Hence it can conclude that TOF based pellets significantly impact the micronutrient availability in soil compared to the FYM based and control treatment.

Keywords: Thermochemical organic fertilizer; pellets; nutrient release; micronutrients; manganese.

1. INTRODUCTION

Micronutrients are essential for plant metabolism, which in turn affects crop production in addition to crop plant growth and development. These micronutrients function as co-factors for a variety of enzymes involved in the metabolism of different organic compounds, including lipids, proteins, carbohydrates, and nucleic acids [1]. Regardless of the source, organic manures are preferable for providing soil with vital nutrients and various micronutrients, including Fe, S, Mo, Zn, and Cu. Upon mineralization, soil organic matter releases considerable amount of phosphorus, sulphur, nitrogen, and a small amount of micronutrients [2,3]. The binding of metals to organic matter may cause a decrease in the concentration of free cations in soil solution; however, the dissolution of these organo-metallic complexes increases the Phyto-availability of metals at the root-rhizosphere interface by raising the concentration of total dissolved ions. This concentration is dependent on the mobility of the metal-dissolved organic carbon (DOC) complexes and their dissociation kinetics. The increase in root-accessible forms of zinc and iron is mostly caused by chelating these elements with organic matter, which also keeps insoluble forms like carbonates and oxides from forming in the soil [4]. Numerous researches have shown the direct and indirect effects of soil organic matter (SOM) on soil availability of micro nutrients and plant absorption [5,6,7]. Dhaliwal and Walia [8] observed that incorporation of manures besides giving the macronutrients it also boosts the availability of the micronutrients like Zn, Cu, Fe, and Mn. Integrated application of organic and inorganic fertilizer is considered to be sustainable then the sole use of inorganic fertilizers. The application of mineral fertilizers and organic manures over time has a substantial impact on soil characteristics, including pH, organic matter contents, and accessible forms of macronutrients, which impacts micronutrient availability to plants [9,10]. Furthermore, adding

organic materials like farmyard manure (FYM) to a balanced fertilization strategy greatly boosts the soil's availability of micronutrients [11]. Pellet fertilizer, integrating both organic and inorganic sources of nutrients, has slow-release nutrient capability, can limit leaching losses, and attributes favorable outcomes on soil nutrient levels and overall health [12].

Hence the current study was conducted to assess the micronutrient dynamics in soil under integrated thermochemical organic fertilizer pellets.

2. MATERIALS AND METHODS

2.1 Soil Incubation Study

A soil incubation study was carried out in the Department of Soil Science and Agricultural Chemistry, to assess dynamics of micronutrient release. Soil Pots were filled with 5 kg air-dried, 2 mm sieved soil. The study was laid in Completely Randomized Design with 5 treatments and 4 replications. The collected soil was sandy clay loam in texture with a moderately acidic pH of 5.87. The Fe, Mn, Zn, Cu and B content of soil was 10.14 mg kg⁻¹, 8.59 mg kg⁻¹, 1.90 mg kg⁻¹, 2.50 mg kg⁻¹ and 0.30 mg kg⁻¹, respectively. Treatments were, T₁: Pellet formulation 1; T₂: Pellet formulation 2; T₃: Pellet formulation 3; T₄: FYM+NPK (RDF as per POP) and T₅: Control (Soil alone). Three distinct pellet formulations P₁, P₂ & P₃ were used in study based on thermochemical organic fertilizer (TOF). P₁, P₂ & P₃ were prepared by blending of TOF with 125%, 100% and 75% of NPK nutrients, respectively based on blanket recommendation for tomato by package of practices of KAU [13]. TOF was mixed mechanically with NPK nutrient sources, homogenized and pelletized using a pelleting machine equipped with cylindrical press channels. The pellets were mixed with soil and incubated for 75 days and the pots were maintained at field capacity throughout the study

period. Soil samples were drawn at 15th, 45th and 75th day. Samples were analysed for available Fe, Mn, Zn and Cu, by 0.1 N HCl extraction and atomic absorption spectrophotometry [14] and B by hot water extraction and Azomethine – H method [15].

2.2 Statistical Analysis

A completely randomized design (CRD) was used for the soil incubation experiment. Critical differences were computed with a 5% significance level in order to facilitate mean comparisons if the effects were significant. Data analysis was done using the R-package grapesAgri1 [16].

3. RESULTS AND DISCUSSION

3.1 Micronutrient Dynamics in the Soil

The Fe and Mn content of the soil showed an increasing trend in all the treatments throughout the incubation period (Table 1). Highest available Fe was recorded in T₁ on 75th day (21.93 mg kg⁻¹) followed by T₂ (21.82 mg kg⁻¹). FYM based treatment showed highest Fe content of 20.70 mg kg⁻¹ on 45th day, unlike treatments amended with pellets which maintained highest availability of Fe throughout

the period. Also control showed an increasing trend in Fe release.

Manganese also exhibits a steady and progressive rate of release throughout, in all the treatments (Table 1). T₁ displayed the highest available Mn (12.76 mg kg⁻¹) on 75th day, followed by T₄ (12.13 mg kg⁻¹) on 75th day (Table 1). Similar to Fe release, control showed similar pattern in Mn release.

The micronutrients Fe and Mn, exhibited a steady and progressive rate of mineralization all throughout the incubation period (Fig. 1, Fig. 2). Citrate, oxalate, malate, malonate, and succinate are all capable of acting as chelators for iron, copper, and zinc, according to Cline et al. [17]. Solubility of plant-available Fe, Mn, Zn, and Cu rises with decreasing soil pH [18]. The genesis of chelating compounds with organic components may be responsible for the steady rise in Fe content over time, as this leads to a slow release throughout the processes of the decomposition and mineralization [19]. The gradual, prolonged release of Fe from organic additions may result from the formation of organic complexes with higher stability with Fe [20] that will likely break down more slowly. The levels of exchangeable Fe and Mn in the soil treated with cow manure increased, according to Rostami and Ahangar [21].

Table 1. Effects of integrated thermochemical organic fertilizer (TOF) pellets on Fe (mg kg⁻¹) and Mn (mg kg⁻¹) at various days of incubation

Treatment details	Fe (mg kg ⁻¹)			Mn (mg kg ⁻¹)		
	Days of incubation			Days of incubation		
	15 Days	45 Days	75 Days	15 Days	45 Days	75 Days
P1	19.65	19.80	21.93	6.50	8.81	12.76
P2	19.33	19.50	21.82	6.52	7.68	10.89
P3	16.27	19.00	21.23	6.34	7.55	11.15
FYM+NPK	13.24	20.70	21.75	8.28	8.28	12.13
Control	12.80	17.24	17.27	5.55	6.73	9.28
SEm (±)	0.719	0.653	0.607	0.156	0.251	0.099
CD (0.05)	2.168	1.970	1.828	0.471	0.756	0.298
	B (mg kg ⁻¹)					
	Days of incubation					
	15 Days	45 Days	75 Days			
P1	0.560	0.570	0.490			
P2	0.545	0.525	0.467			
P3	0.518	0.515	0.440			
FYM+NPK	0.518	0.488	0.440			
Control	0.470	0.470	0.402			
SEm (±)	0.012	0.014	0.005			
CD (0.05)	0.037	0.042	0.016			




Fig. 1. Effect of integrated TOF pellets on soil available Fe




Fig. 2. Effect of integrated TOF pellets on soil available Mn

The rise in Mn release that was observed is consistent with the findings of a study conducted by Radhakrishnan and Suja [22], which showed that Mn and Fe availability increased over many days. The increase in Mn availability from TOF-based treatments up to 12 weeks of incubation was also observed by Ajayan [23]. According to Leno and Sudharmaidevi [5] thermochemical organic fertilizer performed better in the soil at supplying plant-available micronutrients than conventional farmyard manure-based fertilizers.

Zn availability exhibited a slight decline on day 45 and it increased subsequently. The highest value of available zinc was recorded in T₁ on day 15 (5.32 mg kg⁻¹) and day 75 (5.28 mg kg⁻¹) followed by T₂ (5.07 mg kg⁻¹) on day 15 and T₁ (5.06 mg kg⁻¹) on day 45 (Table 2). Whereas, the release of copper showed a decreasing trend after the 15th day of incubation. On day 15 the

highest available copper was recorded in T₁ (5.61 mg kg⁻¹) which was significantly on par with T₂ (5.40 mg kg⁻¹) followed by T₃ (5.11 mg kg⁻¹) (Table 2).

The release pattern of B is similar to the pattern of Cu. The highest available boron content was recorded in in T₁ on days 45 (0.570 mg kg⁻¹) and 15 (0.560 mg kg⁻¹) followed by T₂ on days 15 (0.545 mg kg⁻¹) and 45 (0.525 mg kg⁻¹) (Table 1).

A consistent decrease and increase pattern in Zn release was observed from 15th day (Fig. 3). The decomposition of OM facilitates an increase in the availability of Zn. This process may be attributed to the presence of various functional groups, such as phenolic, carboxyl, and amino groups, which interact with labile organic compounds and create strong chelation, as stated by Fuente et al. [24]. Behera and

Thiyagarajan [25] reported similar results. However, the availability of Zn was reduced because it formed strong bonds with inorganic soil colloids, primarily Fe and Al oxides, through specific adsorption [21].

Cu release increased on initial days and thereby kept on decreasing from 15th day (Fig. 4). The increase soil Cu content following the application of OM can be attributed to the adsorption of Cu²⁺ ions onto the OM. The sorption and complexation processes with OM significantly influence the availability of Cu, as copper tends to form strong inner-sphere complexes with SOM [26]. Additionally, the application of organic manures increases the organic-bound fraction of Cu, while simultaneously reducing its soluble and extractable fractions [27] which eventually decrease the Cu availability for plant absorption.

Despite of high mineralization of B from TOF based pellets, it was deficient in the soil and exhibited a decreasing trend during incubation

(Fig. 5). According to Tlili et al. [28], organic matter is one of the main sources of B in acid soils since relatively little B adsorption on the mineral fraction happens at low pH values. Soil organic matter serves as a significant source of B reserves. Jacoby et al. [29] have demonstrated that organic fertilization has a beneficial effect on B absorption and that B is more prevalent in soils with organic matter than in the mineral component.

Kamaraj et al. [30] found that fortifying organic manures with micronutrients provides greater benefits due to the chelation of these nutrients by organic matter, facilitates a gradual release tailored to crop requirements. The progressive increase observed in the control treatment, which did not receive any amendments, can be attributed to the dissolution of inherent nutrients. This phenomenon occurred because the soil was maintained at field capacity for the entire duration of the study period.

Table 2. Effects of integrated thermochemical organic fertilizer (TOF) pellets on Zn (mg kg⁻¹), Cu (mg kg⁻¹) and B (mg kg⁻¹) at various days of incubation

Treatment details	Zn (mg kg ⁻¹)			Cu (mg kg ⁻¹)		
	Days of incubation			Days of incubation		
	15 Days	45 Days	75 Days	15 Days	45 Days	75 Days
P1	5.32	5.06	5.28	5.61	4.02	4.28
P2	5.07	4.52	4.63	5.40	3.92	3.90
P3	4.46	4.00	4.19	5.11	3.71	3.71
FYM+NPK	3.88	3.33	3.89	4.10	2.93	3.80
Control	3.48	3.05	3.48	3.45	2.78	3.10
SEm (±)	0.078	0.097	0.295	0.153	0.119	0.151
CD (0.05)	0.234	0.294	0.890	0.461	0.358	0.457




Fig. 3. Effect of integrated TOF pellets on soil available Zn




Fig. 4. Effect of integrated TOF pellets on soil available Cu




Fig. 5. Effect of integrated TOF pellets on soil available B

4. CONCLUSION

The study can be concluded with the fact that TOF based pellets significantly impact the micronutrient dynamics in soil. Fe, Mn and Zn exhibited a gradual and progressive release pattern throughout the incubation period. Whereas Cu and B availability exhibited a decreasing pattern from 15th day of incubation. The TPF based pellets generally maintained higher micronutrient availability in the course of the incubation study. P₁ (TOF + NPK @ 125%) comes out to be superior in maintaining the micronutrient availability during the study.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

Authors are thankful to Department of Soil Science and Agricultural Chemistry, College of Agriculture, Kerala Agricultural University, Vellayani, Trivandrum, India for providing all facilities required during course of study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Barker AV, Pilbeam, DJ. editors. Handbook of plant nutrition. 2nd ed. CRC press; 2015.
- Rahman MH, Islam MR, Jahiruddin M, Puteh AB, Mondal MMA. Influence of organic matter on nitrogen mineralization pattern in soils under different moisture regimes. *International Journal of Agriculture & Biology*. 2013;15(1):55–61.
- Amanullah MM. N release pattern in poultry manured soil. *Journal of Applied Sciences Research*. 2007;3:1094–1096.
- Schulin R, Khoshgoftarmansh A, Afyuni M, Nowack B, Frossard E. Effects of soil management on zinc uptake and its bioavailability in plants. In: Banuelos, G., Lin, Z. editors. *Development and Uses of Biofortified Agricultural Products*. CRC Press, Boca Raton, USA: 2009;95–114.
- Leno N, Sudharmaidevi CR. Micronutrient dynamics on addition of a rapid organic fertilizer produced from degradable waste in banana. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(01):1095-1102.
- Ramesha GK, Leno N, Radhika NS. Linking root phenomics, nutrient acquisition and utilisation in amaranthus with thermochemical organic fertilizer from biowaste. *Rhizosphere*. 2021;20:100426.
- Sarangi SS, Sarma I, Gogoi S, Barooah A. Effect of organic nutrients and bio fertilizers on soil parameters and nutritional content of amaranth. *International Journal of Environment and Climate Change*. 2023;13(11):4326-4330.
- Dhaliwal SS, Walia SS. Integrated nutrient management for sustaining maximum productivity of rice-wheat system under Punjab conditions. *Journal of Research, PAU*. 2008;45:12–16.
- Rutkowska B, Szulc W, Sosulski T, Stępień W. Soil micronutrient availability to crops affected by long-term inorganic and organic fertilizer applications. *Plant Soil and Environment*. 2014;60:198–203.
- Brempong MB, Addo-Danso A. Improving soil fertility withorganic fertilizers. In: Turan M, Yildirim E, (eds.) *New generation of organic fertilizers*. Rijeka, Croatia: Intech Open. 2022;1–13.
- Singh V, Rai HK, Jain P, Dwivedi BS, Upadhyay AK, Agrawal SB, Behera SK, Singh RB. Effect of Inorganic and Integrated Long-term Nutrient Management on DTPA-extractable Micronutrients in a Vertisol under Soybean-wheat Cropping System across the Soil Depth. *International Journal of Environment and Climate Change*. 2023; 13(7):471-8.
- Kim P, Hensley D, Labbé N. Nutrient release from switchgrass-derived biochar pellets embedded with fertilizers. *Geoderma*. 2014;232:341-51.
- KAU (Kerala Agricultural University) Package of Practices Recommendations: Crops (15th Ed.). Kerala Agricultural University, Thrissur. 2011;360.
- Sims JT, Johnson GV. Micronutrient soil tests. *Micronutrients in Agriculture*. 1991; 42(4): 427-476.
- Gupta UC. A simplified method for determining hot-water soluble boron in podzol soils. *Soil Science*. 1967; 103 (6): 424-8.
- Gopinath PP, Parsad R, Joseph B, AdarshVS. grapesAgri1: Collection of Shiny Appsfor Data Analysis in Agriculture. *Journal of Open Source Software*. 2021;6 (63): 3437.
- Cline GR, Powell PE, Szanislo PJ, Reid CPP. Comparison of the abilities of hydroxamic, synthetic and other natural organic acids to chelate iron and other ions in nutrient solution. *Soil Science Society of American Journal*. 1982;46(6):1158-1164.
- Brady NC, Weil RR. *The nature and properties of soils*, 13th Ed. Prentice Hall Inc., New Jersey, USA. 2002;960.
- Dhaliwal SS, Naresh RK, Mandal A, Singh R, Dhaliwal, MK. Dynamics and transformations of micronutrients in agricultural soils as influenced by organic matter build-up: A review. *Environmental and Sustainability Indicators*. 2019;1-2: 100007.
- Barber SA. *Soil nutrient bioavailability*, 1st edn. John Wiley & Sons, New Delhi. 2014; 432.
- Rostami GH, Ahangar AG. The effect of cow manure application on the distribution fractions of Fe, Mn and Zn in agricultural soils. *Journal of Agriculture and Veterinary Science*. 2013;6(2): 60-66.

22. Radhakrishnan ARS, Suja G. Nutrient release pattern of organic and inorganic resources used in cassava production (*Manihot esculenta* Crantz). *Journal of Plant Nutrition*. 2019;42(11-12):1301-1315.
23. Ajayan AS. Effect of thermochemical organic fertilizer on soil carbon pools, nutrient dynamics and crop productivity in Ultisols. Ph.D thesis, Kerala Agricultural University, Thrissur. 2021;394.
24. Fuente C, Clemente R, Martinez-Alcala I, Tortosa G, Bernal MP. Impact of fresh and composted solid olive husk and their water-soluble fractions on soil heavy metal fractionation; microbial biomass and plant uptake. *J. Hazardous Mater*. 2011;186 (2-3):1283-1289.
25. Behera PR, Thiyagarajan C. Release kinetics of Zn from soil and growing media mixtures in a laboratory study. *J. Environ. Biol*. 2022;43(3):440-447.
26. Boudesocque S, Guillon E, Aplincourt M, Marceau E, Stievano L. Sorption of Cu (II) onto vineyard soils: macroscopic and spectroscopic investigations. *J. Colloid Interface Sci*. 2007;307(1):40-49
27. Bolan N, Adriano D, Mani S, Khan A. Adsorption, complexation, and phytoavailability of copper as influenced by organic manure. *Environ. Toxicol. Chem*. 2003;22(2):450-456.
28. Tlili A, Dridi I, Attaya R, Gueddari M. Boron characterization, distribution in particle-size fractions, and its adsorption-desorption process in a semiarid Tunisian soil. *Journal of Chemistry*. 2019;(1):1-8.
29. Jacoby R, Peukert M, Succurro A, Koprivova, A, Kopriva S. The role of soil microorganisms in plant mineral nutrition current knowledge and future directions. *Frontiers in Plant Science*. 2017;8:1617.
30. Kamaraj S, Singaravel R, Vembu G. Influence of fortified organic manures on the changes in physicochemical properties and nutrient availability in coastal saline sodic soil for rice. *Asian Journal of Soil Science*. 2010;5(1):60- 64.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/121139>