



Impact of Long Term Tillage and Soil Moisture Regime on the Sulphur, Organic Carbon, DOC, MBC and MBN of Soil under Mung Bean-Wheat and Sorghum-Wheat Cropping System

Deepak Kochar ^{a*} and Rohtas Kumar ^a

^a Department of Soil Science, C.C.S. Haryana Agricultural University, Hisar (Haryana), India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrb/2024/v14i5312>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120653>

Original Research Article

Received: 01/06/2024
Accepted: 02/08/2024
Published: 16/08/2024

ABSTRACT

The field experiment entitled was conducted during 2021-22 on an on-going long term experiment which was started since 2006-07 at Soil Research Farm, Department of Soil Science, CCS HAU, Hisar. The field experiment consisted of three tillage treatments i.e. CT-CT (Conventional tillage in both *kharif* and *rabi* season), ZT-CT (Zero tillage in *kharif* and conventional tillage in *rabi*) and ZT-ZT (Zero tillage in both *kharif* and *rabi* season). The experiment also consisted of two cropping systems MW and SW (Mung bean-wheat and sorghum-wheat) and three soil moisture regimes (IW/CPE= 0.60, 0.75 and 0.90) with three replications. The results of the experiments revealed that the

*Corresponding author: E-mail: deepakkochar9@gmail.com;

Cite as: Kochar, Deepak, and Rohtas Kumar. 2024. "Impact of Long Term Tillage and Soil Moisture Regime on the Sulphur, Organic Carbon, DOC, MBC and MBN of Soil under Mung Bean-Wheat and Sorghum-Wheat Cropping System". *Asian Journal of Research in Biochemistry* 14 (5):62-71. <https://doi.org/10.9734/ajrb/2024/v14i5312>.

sulphur, organic carbon, DOC, MBC and MBN content of the soil were higher under MW than SW cropping system. The highest mean value of sulphur, organic carbon, DOC, MBC and MBN was reported under ZT-ZT followed by ZT-CT and the lowest mean value was under CT-CT under both cropping systems. Similarly, the soil moisture regime M_{0.90} reported the highest mean value of available sulphur, organic carbon, DOC, MBC and MBN content followed by M_{0.75} whereas the lowest mean value was under M_{0.60} under both the cropping system.

Keywords: Zero tillage; cropping system; DOC; MBC; organic carbon.

1. INTRODUCTION

“Mechanical manipulation of soil to provide favourable conditions for proper crop growth is called tillage. Soil tillage consists of breaking the compact surface of earth to a certain depth and loosening the soil mass to enable the roots of the crops to penetrate and spread into the soil. The main objectives of the tillage is to prepare a desirable soil structure for a deep seed bed or a root bed suitable for different types of crops, to control weeds and to minimize soil erosion. Tillage that is deeper and more thorough is classified as primary, and tillage that is shallower and sometimes more selective of location is secondary. Primary tillage such as ploughing tends to produce a rough surface finish, whereas secondary tillage tends to produce a smoother surface finish, such as that required to make a good seedbed for many crops” [1,2,3]. Tillage has been an important part of agriculture history because it achieve many agronomic practices. But the excessive tillage practices affect soil health, crop productivity and environmental quality by affecting soil carbon loss and emission of green house gases [4]. “Therefore, to counteract all these constraints conservation agriculture has been promoted which enhance soil quality related to carbon sequestration that favour soil biology and physical condition” [5,6]. “Conservation agriculture promotes minimum soil disturbance, maintenance of permanent soil cover and diversification of plant species. Conservation agriculture is increasing worldwide due to its economic and environmental benefits for farmers, the environment and society. Economic benefits may arise from lower drought susceptibility due to higher plant-available soil water content resulting in higher yield stability, the saving of labor and fuel and higher economic returns. Ecological benefits include the increase of soil organic carbon (SOC), biotic activity, soil porosity, agro-ecological diversity and less soil erosion and carbon emissions” [7]. “Soil tillage influences the soil chemical characteristics, carbon sequestration and nutrient distributions. Similar to tillage, soil moisture is also an

important parameter which affects the physical, chemical and biological properties of the soil. Soil water plays an important role in maintaining the optimum pH and salinity condition which is essential for plant growth” [8]. Soil water contains nutrients that move into the plant roots when plants take in water. The mineralization and immobilization of nutrients in the soil was controlled by microbes which required an optimum soil moisture. Yang *et al.* [9] “reported that optimum water conditions increase the soil microbial carbon, nitrogen, soil urease activity and soil translocase. Liu *et al.* [10] also reported that adequate soil moisture increase the species and numbers of nitrogen-fixing bacteria, gamma-deformation bacteria and alpha deformation bacterial of the soil”. “Extreme moisture conditions would limit CO₂ release: low soil moisture would limit microbial and root respiration; excessive would block the soil pore and limit O₂ and CO₂ release. Generally, SOC mineralization rate increases with the soil water content” [11]. This study aimed to influence the impact of long term tillage and soil moisture regime on the sulphur, organic carbon, DOC, MBC and MBC of the soil under mung bean-wheat and sorghum-wheat cropping system.

2. MATERIALS AND METHODS

2.1 Experiment Site Characteristics

The field experiment was conducted from the on-going experiment, which was started since 2006-07 at Soil Research Farm, Department of Soil Science, CCS HAU, Hisar. Hisar is the administrative headquarters of Hisar district in the state of Haryana in northwestern India. Hisar is located at 29.09°N 75.43°E in western Haryana. It has an average elevation of 215 m above mean sea level. The soil of the experimental site was sandy loam in texture, slightly alkaline, low in organic carbon, low in available nitrogen, medium in available phosphorus and high in available potassium (Phogat, 2019). The main characteristics of the climate of the experiment site are dryness,

extremes of temperature, and scanty rainfall. The maximum daytime temperature during the summer varies between 40 and 46 °C. During winter, it ranges between 1.5 °C and 4 °C. Relative humidity varies from 5 to 100%. The average annual rainfall is around 429 mm, most of which occurs during July and August. Dew is observed in December and January. Hot winds, locally known as loo, are strong and frequent from May to July.

2.2 Treatments and Experimental Design

“The experiment was carried out in two main plots treatments viz. (i) Mungbean-wheat and (ii) Sorghum-wheat cropping system. These main plots were split in to sub plots receiving three different tillage practices i.e (i) Conventional tillage in both *kharif* & *rabi* seasons (CT-CT), (ii) conventional tillage in *kharif* & zero tillage in *rabi* seasons (CT-ZT) and, (iii) zero tillage in both *kharif* & *rabi* seasons” (ZT-ZT). [12] These sub-plots were further split into sub-sub plots receiving different soil moisture regimes viz. IW/CPE =0.60, 0.75 and 0.90 to each sub-plots. The experiment was layout in split plot design and size of each plot was 35m x 8m. Replications were three of each treatments and in conventional tillage field was ploughed during both season. In second treatment plots, the fields were ploughed during *kharif* only and no tillage was done during *rabi* season. In zero tillage plots, no tillage was done during both the seasons. The residues of the preceding crop were removed in conventional tillage. In zero tillage practice plots, the crop was harvested and no-tillage was done for preparation of the seedbed for the succeeding crop, and zero till machine was used for sowing the crop.

2.3 Soil Sampling and Measurement

The soil samples were collected from 0-15 cm depth after the harvest of the crop in the end of the *kharif* season. Three samples from each of the plots were collected with the help of augur and khurpi and made a composite sample by mixing these samples. These composite sample were brought to the laboratory in the sampling bag, dried in shade, grind, sieved (2mm) and finally stored in the sampling bag with proper labeling. Available sulphur was determined was estimated turbidimetric method developed by Chesnin and Yien, [13]. Dissolved organic carbon was determined by dichromate acid oxidation method [14]. Total soil organic carbon was determined using Walkley and Black [15] wet digestion method. Microbial biomass carbon

was determined by the chloroform (CHCl₃) fumigation method proposed by Vance et al., [16]. Microbial biomass nitrogen in soil samples was estimated as per the method of Brookes et. al. [17].

2.4 Statistical Analysis

The data generated in both laboratory and field experiments were statistically analysed as per the design of the experiment. Treatment effects were compared with CD (0.05). Statistical analysis was done in consultation with Department of Mathematics and Statistics, CCS Haryana Agricultural University, Hisar.

3. RESULTS

3.1 Available Sulphur

The sulphur content under mung bean-wheat cropping system was significantly higher than sorghum-wheat (Table. 1). The effect of tillage practices on available sulphur was significant and highest mean value of available sulphur was 28.00 and 20.44 kg ha⁻¹ under ZT-ZT. The available sulphur in this treatment was statistically at par with ZT-CT and significantly different from CT-CT in mung bean-wheat cropping system but in the sorghum-wheat cropping system it was statistically at par with both CT-CT and ZT-ZT treatment. Also, the sulphur content under ZT-CT treatment had mean value of 26.87 and 19.86 kg ha⁻¹ which was also statistically at par with CT-CT treatment. The lowest value of available sulphur was 24.99 and 18.58 kg ha⁻¹ under CT-CT treatment. The effect of soil moisture regimes was found significant on available sulphur under both cropping systems. The highest mean value of available sulphur was 29.47 and 20.44 kg ha⁻¹ under soil moisture regime M_{0.90} which was significantly higher than M_{0.75} and M_{0.60}. Similarly, the mean value of available sulphur under M_{0.75} was 26.28 and 19.65 kg ha⁻¹ which was significantly higher than M_{0.60}. The lowest mean value of available sulphur was 25.12 and 18.79 kg ha⁻¹ in the plots under M_{0.60}. However, the combine effect of cropping system and tillage; cropping system and soil moisture regimes; tillage and soil moisture regimes; cropping system, tillage and soil moisture regimes on available sulphur was found non-significant.

3.2 Dissolved Organic Carbon (DOC)

The DOC was found significantly higher under the sorghum-wheat cropping system then mung

bean-wheat (Table. 2). The highest DOC mean value (0.54 and 0.48 g kg⁻¹) under both cropping system was recorded under ZT-ZT treatment and lowest (0.49 and 0.41 g kg⁻¹) was under CT-CT treatment. The mean value of DOC under ZT-ZT was found significantly higher than ZT-CT and CT-CT under both cropping system. Also, the mean value of DOC under the ZT-CT tillage treatment was 0.52 and 0.44 g kg⁻¹ which was also significantly higher than the CT-CT treatment. The present study revealed that the DOC content of the plots under different soil moisture regime was also significantly affected and increased as the soil moisture regimes increased. The highest mean value of DOC was 0.54 and 0.47 g ka⁻¹ under M_{0.90}. This mean value of DOC was found significantly higher than the M_{0.75} and M_{0.60} under both cropping system. Similarly, the mean value of DOC (0.51 and 0.44 g kg⁻¹) which was under M_{0.75} soil moisture regimes was also significantly higher than M_{0.60}. The lowest mean value of DOC was 0.50 and 0.42 g kg⁻¹ under soil moisture regime M_{0.60}. However, the interactive effect of cropping system and tillage; cropping system and soil moisture regimes was found significant whereas tillage and soil moisture regimes; cropping system, tillage and soil moisture regimes were found non-significant.

3.3 Soil Organic Carbon

The organic carbon was reported significantly higher under mung bean-wheat than sorghum-wheat cropping system (Table. 3). However, the soil under ZT-ZT recorded the highest (0.77 and 0.75 %) mean value of organic carbon which was significantly higher than ZT-CT and CT-CT under both cropping system. The lowest (0.69 and 0.67 %) mean value of organic carbon was observed under CT-CT under both cropping system. Also, the soil under ZT-CT mean value of organic carbon was 0.74 and 0.72 % under both cropping system respectively which was significantly higher than CT-CT. Similarly, long term effect of the moisture regime also significantly affects the organic carbon of soil. It was observed that the organic carbon of soil increased as soil moisture regimes increased. The highest mean value of organic carbon was observed under under M_{0.90} (0.76 and 0.74 %) under both system respectively. The organic carbon in the soil under M_{0.90} was found significantly higher than M_{0.75} and M_{0.60}. Also, the soil under M_{0.75} moisture regimes had mean value of soil organic carbon 0.73 and 0.71 % in both cropping system which was significantly

higher than M_{0.60} whereas the lowest mean value of soil organic carbon (0.71 and 0.69 %) was observed under M_{0.60}. However, the combined effect of cropping system and tillage; cropping system and soil moisture regimes; cropping system, tillage and soil moisture regimes was found non-significant.

3.4 Microbial Biomass Carbon (MBC)

The MBC of the soil was significantly affected by the cropping system and it was higher under mung bean-wheat cropping system as compared to sorghum-wheat (Table. 4). The mean MBC which was under ZT-ZT were 838 and 817 mg kg⁻¹ which was found significantly higher than ZT-CT and CT-CT under both cropping system. Similarly, the mean MBC (761 and 738 mg kg⁻¹) under ZT-CT was also significantly higher than which was under CT-CT under both cropping system. The lowest mean value of MBC was 681 and 662 mg kg⁻¹ which was reported under CT-CT tillage treatment under both cropping system. This data also show that the MBC was also significantly affected by the different soil moisture regimes. It was observed that the mean value of the MBC was increased as the soil moisture regime increased. The soil which was under M_{0.90} gave the highest mean value of MBC (868 and 851 mg kg⁻¹) under both cropping system which was found significantly higher than the mean MBC of other two moisture regimes. The mean value of MBC which was under M_{0.75} was 748 and 735 mg kg⁻¹ under both the cropping system respectively. These mean value of MBC under M_{0.75} soil moisture regimes was significantly higher than M_{0.60} under both cropping system whereas the lowest (663 and 630 mg kg⁻¹) mean value of MBC was reported under M_{0.60} soil moisture regime under both cropping system. The cropping pattern and tillage; cropping pattern and soil moisture regimes; tillage and soil moisture regimes; cropping pattern, tillage and soil moisture regimes also significantly effect the MBC of the soil. In both cropping system, the soil under ZT-ZT and M_{0.90} had the highest value of MBC was 919 mg kg⁻¹ which was significantly higher than rest of the treatment whereas the lowest (545 mg kg⁻¹) value was under CT-CT and M_{0.60} which was also significantly lower than rest of the treatment.

3.5 Microbial Biomass Nitrogen

The MBN of the soil of plots was significantly affected by the cropping system and record

higher in mung bean-wheat cropping system than sorghum-wheat (Table. 5). The highest mean value (135.7 and 121.6 mg kg⁻¹) of MBN was observed under ZT-ZT treatment followed by ZT-CT (106.3 and 96.1 mg kg⁻¹) under both the cropping system respectively. The mean value of MBN which was observed in the ZT-ZT was significantly higher than the other two tillage practices whereas ZT-CT was significantly higher than CT-CT under both cropping system. The lowest mean value of MBN was recorded in CT-CT tillage treatment which was 90.5 and 85.4 mg kg⁻¹ under both the cropping system respectively. Present data revealed that the

effect of different soil moisture regimes on the MBN was found significant and was highest in the plots which was under M_{0.90}. It was reported that the mean value of MBN was increased as the value of soil moisture regime increased. The mean value of MBN which was under M_{0.90} was 122.0 and 108.9 mg kg⁻¹ under both cropping system which was found significantly higher than the other two soil moisture regimes. The soil which was under soil moisture regime M_{0.75} had mean values of MBN 103.9 and 96.7 mg kg⁻¹ which was significantly higher than M_{0.60}. The lowest mean value (87.4 and 83.7 mg kg⁻¹) of MBN was recorded in the plots which was under

Table 1. Effect of moisture regimes and tillage on the available sulphur (kg ha⁻¹) of soil under mung bean-wheat and sorghum-wheat cropping system

Moisture Regime (IW/CPE)	Mung bean-Wheat			Mean	Sorghum-Wheat			Mean
	ZT-ZT	ZT-CT	CT-CT		ZT-ZT	ZT-CT	CT-CT	
M _{0.60}	26.49	24.70	24.19	25.12	19.59	19.29	17.50	18.79
M _{0.75}	28.08	26.75	24.00	26.28	20.33	19.77	18.96	19.65
M _{0.90}	30.44	29.18	28.79	29.47	21.51	20.52	19.27	20.44
Mean	28.00	26.87	24.99		20.44	19.86	18.58	

CD (p=0.05) A = 5.31, B = 1.95, C = 0.35, AXB = NS, AXC= NS, BXC= NS, AXBXC=NS
 CT = conventional tillage, ZT = zero tillage, M_{0.60} = moisture regime at IW/CPE=0.60, M_{0.75}= moisture regime at IW/CPE= 0.75, M_{0.90}= moisture regime at IW/CPE=0.90; A= cropping factor, B= tillage factor, C= moisture regime factor

Table 2. Effect of moisture regimes and tillage on the DOC (g kg⁻¹) of soil under mung bean-wheat and sorghum-wheat cropping system

Moisture Regime (IW/CPE)	Mung bean-Wheat			Mean	Sorghum-Wheat			Mean
	ZT-ZT	ZT-CT	CT-CT		ZT-ZT	ZT-CT	CT-CT	
M _{0.60}	0.52	0.51	0.48	0.50	0.45	0.41	0.39	0.42
M _{0.75}	0.54	0.50	0.49	0.51	0.49	0.43	0.41	0.44
M _{0.90}	0.57	0.54	0.51	0.54	0.51	0.48	0.44	0.47
Mean	0.54	0.52	0.49		0.48	0.44	0.41	

CD (p=0.05) A = 0.017, B = 0.007, C = 0.009, AXB = 0.010, AXC= 0.012, BXC= NS, AXBXC=NS
 CT = conventional tillage, ZT = zero tillage, M_{0.60} = moisture regime at IW/CPE=0.60, M_{0.75}= moisture regime at IW/CPE= 0.75, M_{0.90}= moisture regime at IW/CPE=0.90; A= cropping factor, B= tillage factor, C= moisture regime factor

Table 3. Effect of moisture regimes and tillage on the soil organic carbon (%) of soil under mung bean-wheat and sorghum-wheat cropping system. [18]

Moisture Regime (IW/CPE)	Mung bean-Wheat			Mean	Sorghum-Wheat			Mean
	ZT-ZT	ZT-CT	CT-CT		ZT-ZT	ZT-CT	CT-CT	
M _{0.60}	0.74	0.71	0.68	0.71	0.72	0.68	0.64	0.69
M _{0.75}	0.77	0.74	0.69	0.73	0.75	0.71	0.67	0.71
M _{0.90}	0.79	0.77	0.71	0.76	0.77	0.74	0.70	0.74
Mean	0.77	0.74	0.69		0.75	0.72	0.67	

CD (p=0.05) A = 0.010, B = 0.015, AXB = NS, C = 0.015, AXC= NS, BXC= NS, AXBXC=NS
 CT = conventional tillage, ZT = zero tillage, M_{0.60} = moisture regime at IW/CPE=0.60, M_{0.75}= moisture regime at IW/CPE= 0.75, M_{0.90}= moisture regime at IW/CPE=0.90; A= cropping factor, B= tillage factor, C= moisture regime factor

Table 4. Effect of moisture regimes and tillage on the microbial biomass carbon (mg kg⁻¹) of soil under mung bean-wheat and sorghum-wheat cropping system

Moisture Regime (IW/CPE)	Mung bean-Wheat			Mean	Sorghum-Wheat			Mean
	ZT-ZT	ZT-CT	CT-CT		ZT-ZT	ZT-CT	CT-CT	
M _{0.60}	748	672	570	663	713	633	545	630
M _{0.75}	847	736	662	748	835	728	643	735
M _{0.90}	919	874	811	868	903	855	797	851
Mean	838	761	681		817	738	662	

CD (p=0.05) A = 0.16, B = 0.19, C = 0.16, AXB = 0.23, AXC= 0.21, BXC= 0.25, AXBXC=0.29

CT = conventional tillage, ZT = zero tillage, M_{0.60} = moisture regime at IW/CPE=0.60, M_{0.75}= moisture regime at IW/CPE= 0.75, M_{0.90}= moisture regime at IW/CPE=0.90; A= cropping factor, B= tillage factor, C= moisture regime factor

Table 5. Effect of moisture regimes and tillage on the microbial biomass nitrogen (mg kg⁻¹) of soil under mung bean-wheat and sorghum-wheat cropping system

Moisture Regime (IW/CPE)	Mung bean-Wheat			Mean	Sorghum-Wheat			Mean
	ZT-ZT	ZT-CT	CT-CT		ZT-ZT	ZT-CT	CT-CT	
M _{0.60}	96.5	86.7	79.1	87.4	91.6	88.6	71.0	83.7
M _{0.75}	117.8	103.8	90.2	103.9	110.5	93.0	86.8	96.7
M _{0.90}	135.7	128.2	102.2	122.0	121.6	106.8	98.3	108.9
Mean	116.7	106.3	90.5		107.9	96.1	85.4	

CD (p=0.05) A = 3.8, B = 1.9, C = 1.5, AXB = 2.6, AXC= 2.2, BXC= 2.7, AXBXC=3.8

CT = conventional tillage, ZT = zero tillage, M_{0.60} = moisture regime at IW/CPE=0.60, M_{0.75}= moisture regime at IW/CPE= 0.75, M_{0.90}= moisture regime at IW/CPE=0.90; A= cropping factor, B= tillage factor, C= moisture regime factor

soil moisture regime M_{0.60}. Also, the interactive effect of cropping system and tillage; cropping system and soil moisture regimes; tillage and soil moisture regimes; cropping system, tillage and soil moisture regimes were observed significant on the microbial biomass nitrogen of the soil. In both cropping system, the soil under ZT-ZT and M_{0.90} had the highest value of MBN was 135.7 mg kg⁻¹ which was significantly higher than rest of the treatment whereas the lowest (71.0 mg kg⁻¹) value was under CT-CT and M_{0.60} which was also significantly lower than rest of the treatment.

4. DISCUSSION

4.1 Available Sulphur

The available sulphur content of the soil under long term tillage treatment and soil moisture regimes was higher under zero tillage than conventional tillage under both the cropping system. The available sulphur content of the soil was significantly higher under mung bean-wheat cropping system than sorghum-wheat. The available sulphur content of the soil was highest under ZT-ZT followed by ZT-CT and lowest

under CT-CT under both the cropping system. The higher mean value of available sulphur under zero tillage was might be due to higher organic matter which decompose and increase the available sulphur status of the soil. Alam et al. [19] and Shiwakoti et al. [20] reported higher content of available sulphur under zero tillage than conventional tillage. Also, the higher root mass and crop residue under zero tillage increase the sulphur mineralization microbes population which enhance the sulphur content of the soil. Increase in the microbial population under no tillage was also reported by He et al. [21] and Srour et al. [22] in their long term tillage experiments. In the present study, the available sulphur content in the plots increased as the soil moisture regimes increased under both the cropping system. The highest mean value of available sulphur was observed under M_{0.90} followed by M_{0.75} whereas the lowest mean value was under M_{0.60} under both the cropping system. The increase in the availability of sulphur with increase in the moisture content of the soil may be due to high microbial activities and higher mineralization of organic sulphur which increase its availability. Hassan [23] reported higher mineralization and higher

sulphur content at higher soil moisture content. Enhancement in the enzyme and microbial activities with increase in soil moisture content reported by Chen et al. [24] which enhance the mineralization of organic matter which increase the available sulphur content of soil.

4.2 Dissolved Organic Carbon (DOC)

The dissolved organic carbon content of the soil under long term tillage and soil moisture regimes was significantly higher under zero tillage than conventional tillage under both the cropping system. The dissolved carbon content of the soil was higher under mung bean-wheat cropping system than sorghum-wheat cropping system. The higher mean value of dissolved organic carbon under mung bean-wheat cropping system might be due to more crop residue and root biomass with low C:N ration which decompose easily and increases the carbon content. Higher dissolved organic carbon under legume based cropping system was reported by Li et al. [25]. The highest mean value of dissolved organic carbon was observed under ZT-ZT followed by ZT-CT and the lowest mean value was under CT-CT under both the cropping system. The higher value of dissolved organic carbon under zero tillage might be due to accumulation of more crop residue which increase the dissolved organic carbon of soil on decomposition. Increase in the dissolved organic carbon under no till was reported by Dou et al. [26] and Schmidt and Martinez [27]. The dissolved organic carbon content of the soil increases with the increase in the soil moisture regimes under both cropping system. The highest mean value of dissolved organic carbon was observed under $M_{0.90}$ followed by $M_{0.75}$ whereas the lowest mean value was under $M_{0.60}$ under both cropping system. The increase in dissolve organic carbon in soil with increase in soil moisture might be due to higher microbial biomass which decompose the soil organic matter and enhance the carbon status of soil. The positive influence of soil moisture on the microbial biomass was reported by Curtin et al. [28] and Bhanwaria et al. [29]. Yun et al. [11] reported an increases in the dissolve organic carbon of the soil with increase in soil moisture.

4.3 Soil Organic Carbon

The organic carbon content of the soil was significantly higher under mung bean-wheat cropping system than sorghum-wheat which might be due to narrow C:N ratio of the legumes

which decompose easily and increase the SOC content of the soil. Cogreves et al. [30] also reported higher value of SOC under legumes crop under different cropping systems in their findings. The highest mean value of organic carbon was under ZT-ZT followed by ZT-CT whereas the lowest mean value was under CT-CT under both the cropping system. The value of organic carbon was significantly higher under ZT-ZT might be due to less disturbance of soil which increase the organic matter content as well as root mass of the soil. The lower value of organic carbon under CT-CT was due to fast oxidation of the organic matter of the soil which decrease the organic carbon of the soil. Kahlon and Gurpreet et al. [4] and Neugschwandtner [7] reported higher value of organic carbon of the soil under zero tillage as compared to conventional tillage in their field experiments. Also, under different soil moisture regimes the organic carbon was highest under $M_{0.90}$ whereas the lowest value was under $M_{0.60}$ under both the cropping system. The increase in the soil organic carbon with increase in soil moisture regimes might be due to favourable conditions for the microbes to decompose the organic matter which increase the soil organic carbon. Yun et al. [11], and Fang et al. [31] also noticed an increase in the soil organic carbon with increase in the soil moisture content in their field experiments.

4.4 Microbial Biomass Carbon and Nitrogen (MBC and MBN)

The microbial biomass carbon and nitrogen was significantly affected by the cropping system, tillage and soil moisture regimes. The microbial biomass carbon and nitrogen was higher in the soil which was under zero tillage as compared to conventional tillage. The microbial biomass carbon and nitrogen of the soil was higher under mung bean-wheat cropping system than sorghum-wheat cropping system. Higher microbial biomass carbon in the soil under legume based cropping system was reported by Li et al. [25]. The highest mean value of microbial biomass carbon and nitrogen was observed under ZT-ZT followed by ZT-CT and minimum was under CT-CT under both the cropping system. The higher mean value of microbial biomass under zero tillage may be due to more crop residue and less disturbance in the soil which increase the microbial population and provide a steady source of carbon to the microbes which enhanced the microbial biomass carbon and nitrogen of the soil. These results

are similar to the findings of Balota et al. [32], Bausenwein et al. [33] and Yeboah et al. [34] which indicated no tillage generally increase the microbial biomass carbon and nitrogen of soil. The microbial biomass carbon and nitrogen of the soil increase with increase in the soil moisture regimes under both the cropping system because high moisture create favourable condition for the microbial activity. Similar results are found by Bhanwaria et al. [29] and Tuo et al. [35].

5. CONCLUSION

The results of the present experiment concluded that the long term zero tillage and optimum soil moisture regimes improves the chemical properties of the soil as compared to conventional tillage under mung bean-wheat and sorghum-wheat cropping system. The results also show that the legume based cropping system is better than non-legume based cropping system. The results also concluded that long term zero tillage and optimum soil moisture regime increases the available sulphur, organic carbon, DOC, MBC and MBN of the soil as compared with conventional tillage under mung bean-wheat and sorghum-wheat cropping system.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar R, Naresh RK, La LM, Sachan DK, Mahajan NC, Singh S, Singh O, Kumar R, Chaudhary V. Can conservation tillage and residue management effects on sensitivity of labile soil organic carbon fractions and soil organic carbon stocks in sub-tropical ecosystems: A review. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(11):2063-89.
2. Naresh RK, Kumar A, Bhaskar S, Dhaliwal SS, Kumar S, Kumar S, Gupta RK. Organic matter fractions and soil carbon sequestration after 15-years of integrated nutrient management and tillage systems in an annual double cropping system in northern India. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(6):670-83.
3. Priyadarshi R, Kumari A. Effect of conservation tillage and crop rotations on soil organic carbon fractions and soil microbial biomass carbon: A review. *IJCS*. 2019;7(6):1571-8.
4. Kahlon MS, Gurpreet S. Effect of tillage practices on soil physico-chemical characteristics and wheat straw yield. *International Journal of Agricultural Sciences*. 2014;4(10): 289.
5. Bronick CJ, and Lal R. Soil structure and management: a review. *Geoderma*. 2005; 124(1-2): 3-22.
6. Muñoz A, López-Piñero A, Ramírez M. Soil quality attributes of conservation management regimes in a semi-arid region of south western Spain. *Soil and Tillage Research*. 2007;95(1-2):255-265.
7. Neugschwandtner RW, Liebhard P, Kaul HP, Wagentristsl H. Soil chemical properties as affected by tillage and crop rotation in a long-term field experiment; 2014.
8. Chang T, Zhang Y, Zhang Z, Shao X, Wang W, Zhang J, Yang X, Xu H. Effects of irrigation regimes on soil NO₃--N, electrical conductivity and crop yield in plastic greenhouse. *International Journal of Agricultural and Biological Engineering*. 2019;12(1): 109-115.
9. Yang XY, Jiang DH, Yang GR. Effects of water and fertilizer integration on soil microbial biomass carbon, nitrogen and enzyme activities in sugarcane. *Chin. J. Soil Sci*. 2018;49(4): 889–896.
10. Liu ZX, Liu P, Jia XC, et al. Effects of irrigation and fertilization on soil microbial properties in summer maize field. *Chin. J. Appl. Ecol*. 2015;26 (1):113–121.
11. Yun J, Chen X, Liu S, Zhang W. Effects of temperature and moisture on soil organic carbon mineralization. In *IOP conference series: materials science and engineering*. 2019;562(1):012085. IOP Publishing.
12. Mamta Phogat, Rita Dahiya, Sangwan PS, Vishal Goyal, Kakraliya SK, Santhosh Kumar M. Effect of Long Term Zero Tillage and Different Moisture Regimes on Soil Organic Carbon and its Fractions in

- Legume based Cropping Systems of North-western Indo-Gangetic Plains Int.J.Curr.Microbiol.App.Sci. 2021;10(03): 674-693.
13. Chesnin L, Yien CH. Turbidimetric Determination of Available Sulphates. Soil Science Society of America Journal. 1950;15:149-151.
 14. Ciavatta C, Vittori Antisari L, Sequi P. Determination of organic carbon in soils and fertilizers. Communications in Soil Science. 1989;20:1-90.
 15. Walkley A, Black I.A. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. Soil Science. 1934;37: 29-38.
 16. Vance ED, Brookes PC, Jenkinson DS. An extraction method for measuring soil microbial biomass C. Soil biology and Biochemistry. 1987;19(6):703-707.
 17. Brookes PC, Landman A, Pruden G, Jenkinson DS. Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. Soil Biology & Biochemistry. 1985;17: 837-842.
 18. Kochar D, Kumar R. Effect of Long Term Tillage and Soil Moisture Regime on the Chemical Properties of Soil under Mung Bean-Wheat and Sorghum-Wheat Cropping System. International Journal of Plant & Soil Science. 2024;36(7):797-808.
 19. Alam MK, Islam MM, Salahin N, Hasanuzzaman M. Effect of tillage practices on soil properties and crop productivity in wheat-mungbean-rice cropping system under subtropical climatic conditions. The scientific world journal; 2014.
 20. Shiwakoti S, Zheljaskov VD, Gollany HT, Xing B, Kleber M.. Micronutrient concentrations in soil and wheat decline by long-term tillage and winter wheat-pea rotation. Agronomy. 2019;9(7):359.
 21. He JZ, Shen JP, Zhang LM, Zhu YG, Zheng YM, Xu MG, Di H. Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization practices. Environmental microbiology. 2007;9(9):2364-2374.
 22. Srour AY, Ammar HA, Subedi A, Pimentel M, Cook RL, Bond J, Fakhoury AM. Microbial communities associated with long-term tillage and fertility treatments in a corn-soybean cropping system. Frontiers in Microbiology. 2020; 11: 522658.
 23. Hassan MM. Mineralization rate of sulphur in soils of new alluvial zone of West Bengal under regimes of moisture content. J Pharm Innov. 2022;11(4):39-43.
 24. Chen G, Lu Q, Bai J, Wen L, Zhang G, Wang W, Wang C, Liu Z. Organic sulfur mineralization in surface soils from coastal wetlands with different flooding periods affected by the flow-sediment regulation in the Yellow River Delta, China. Catena. 2022;215: 106343.
 25. Li G, Tang X, Hou Q, Li T, Xie H, Lu Z, Zhang T, Liao Y, Wen X. Response of soil organic carbon fractions to legume incorporation into cropping system and the factors affecting it: A global meta-analysis. Agriculture, Ecosystems & Environment. 2023;342: 108231.
 26. Dou F, Wright AL, Hons FM. Dissolved and soil organic carbon after long-term conventional and no-tillage sorghum cropping. Communications in soil science and plant analysis. 2008;39(5-6):667-679.
 27. Schmidt MP, Martínez CE. The influence of tillage on dissolved organic matter dynamics in a Mid-Atlantic agroecosystem. Geoderma. 2019;344: 63-73.
 28. Curtin D, Beare MH, Hernandez-Ramirez G. Temperature and moisture effects on microbial biomass and soil organic matter mineralization. Soil Science Society of America Journal. 2012;76(6):2055-2067.
 29. Bhanwaria R, Singh B, Musarella CM. Effect of organic manure and moisture regimes on soil physiochemical properties, microbial biomass C_{mic}: N_{mic}: P_{mic} turnover and yield of mustard grains in arid climate. Plants. 2022;11(6):722.
 30. Congreves KA, Hayes A, Verhallen EA, Van Eerd LL. Long-term impact of tillage and crop rotation on soil health at four temperate agroecosystems. Soil and Tillage Research. 2015;152:17-28.
 31. Fang X, Zhu YL, Liu JD, Lin XP, Sun HZ, Tang XH, Hu YL, Huang YP, Yi ZG. Effects of moisture and temperature on soil organic carbon decomposition along a vegetation restoration gradient of subtropical China. Forests. 2022;13(4): 578.

32. Balota EL, Colozzi Filho A, Andrade DS, Dick RP. Long-term tillage and crop rotation effects on microbial biomass and C and N mineralization in a Brazilian Oxisol. *Soil and Tillage Research*. 2004;77(2):137-145.
33. Bausenwein U, Gattinger A, Langer U, Embacher A, Hartmann HP, Sommer M, Munch JC, Schloter M. Exploring soil microbial communities and soil organic matter: variability and interactions in arable soils under minimum tillage practice. *Applied soil ecology*. 2008;40(1): 67-77.
34. Yeboah S, Zhang R, Cai L, Li L, Xie J, Luo Z, Liu J, Wu J. Tillage effect on soil organic carbon, microbial biomass carbon and crop yield in spring wheat-field pea rotation. *Plant soil environment*. 2016;62(6): 279-285.
35. Tuo Y, Wang Z, Zheng Y, Shi X, Liu X, Ding M, Yang Q. Effect of water and fertilizer regulation on the soil microbial biomass carbon and nitrogen, enzyme activity, and saponin content of *Panax notoginseng*. *Agricultural Water Management*. 2023; 278: 108145.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/120653>