



Advances in Micropropagation Techniques for Aquascaping Plants: A Comprehensive Review

Greeshma Baby ^{a++*}, M. Rafeekher ^{a#}, Soni, K.B ^{b†},
Priya Kumari I ^{a#}, Reshmi C.R ^{a#}, Sheena.A ^{c#}
and Abdul Rahman M ^{d++}

^a Department of Floriculture and Landscaping, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala, India.

^b Department of Biotechnology, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala, India.

^c Instructional Farm, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala, India.

^d Department of Horticulture, Floriculture and Landscape Architecture, School of Agricultural Sciences, Nagaland University, Medziphema, Nagaland.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/acri/2024/v24i11944>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125306>

Review Article

Received: 18/08/2024

Accepted: 20/10/2024

Published: 24/10/2024

ABSTRACT

Aquascaping is a creative arrangement of aquatic plants, rocks, stones, driftwood, and other elements within an aquarium to create visually pleasing underwater landscapes. It blends principles of design, horticulture, and aquaculture to either mimic natural environments or express artistic

⁺⁺ PhD Scholar;

[#] Assistant Professor;

[†] Professor;

*Corresponding author: Email: greeshmavembenickal@gmail.com;

Cite as: Baby, Greeshma, M. Rafeekher, Soni, K.B, Priya Kumari I, Reshmi C.R, Sheena.A, and Abdul Rahman M. 2024. "Advances in Micropropagation Techniques for Aquascaping Plants: A Comprehensive Review". Archives of Current Research International 24 (11):14-22. <https://doi.org/10.9734/acri/2024/v24i11944>.

creativity. Aquascaping plants are aquatic plants selected and cultivated for their aesthetic appeal and ability to thrive underwater, enhancing the visual beauty of aquarium landscapes. These plants are chosen based on their capacity to flourish when submerged and their role in creating attractive underwater environments. As aquascaping industry is gaining importance, rapid propagation of plants used in it plays pivotal role. Most of the plants are slow growers and their propagation is difficult rendering these plants very expensive. Tissue culture can be a best alternative propagation method in propagation of aquascaping plants. In vitro propagation methods like tissue culture, callus culture is highly relevant and beneficial in the context of aquascaping plants, where precise control over plant growth and health is crucial for creating and maintaining aquascapes

Keywords: Aquascaping; tissue culture; aquariums; In vitro propagation.

1. INTRODUCTION

Aquascaping involves the creative arrangement of aquatic plants, stones, rocks, or driftwood in a visually appealing manner within an aquarium (Martin 2013). It involves creating aesthetically pleasing creations under water using plants, rocks, substrates, and sometimes fish or other aquatic creatures. Aquascaping is a combination of art and horticulture to portray alluring and cordial aquatic scenes, often inspired by seascapes (Baby et al. 2024). Aquascaping, a discipline within the aquarium hobby, has evolved through the contributions of various influential figures and groups over time. Takashi Amano, a notable figure from Japan (1954-2015), is widely recognized as a pioneer in modern aquascaping. He played a focal role in advancing the creation of naturalistic underwater landscapes, promoting layouts inspired by nature and emphasizing principles such as balance, harmony, and the integration of aquatic plants to replicate natural habitats. Tissue culture techniques have offered notable benefits in the efficient and accelerated production of plants (Espinosa-Leal, et al. 2018). Some plants commonly used in aquascaping include Java moss, Anubias, Cryptocoryne, Hygrophila, Ludwigia, Java fern, Amazon sword, Vallisneria, Pogostemon, and Rotala.

Tissue-cultured aquatic plants have seen a sharp increase in commercial interest recently. Tissue culture techniques have been successfully used to cultivate these plants, as numerous research have shown (Jabir et al. 2016). A key factor in the regeneration of shoots is the kind and quantity of growth regulators added to the tissue culture medium (Kunene et al. 2018). One of the key advantages of tissue culture for aquascaping plants is the ability to produce pathogen-free, healthy plantlets (Hussain et al. 2012). This is particularly important in the aquarium setting, where the introduction of diseases or pests can

have devastating effects on the entire ecosystem. Aquascaping plants often prefer tissue culture due to their sterile propagation conditions, which guarantee they are free from pests, diseases, and algae essential for maintaining a healthy aquarium environment. Furthermore, tissue culture plants are consistent in quality and appearance, ensuring uniform growth that aids aquascapers in achieving precise design aesthetics. Additionally, these plants tend to develop strong root systems, facilitating quicker adaptation to underwater conditions compared to traditionally grown plants in pots or soil. In summary, tissue culture plants are valued for their reliability, cleanliness, and ability to thrive, making them a preferred choice for vibrant and thriving aquascaping displays. The demand is limited to specific markets and in five years (1990- 1995) the production in Holland increased from 417,000 to 1,204,000 plants, some of which propagated *in vitro* (Pierik & Ruibing 1997).

2. METHODOLOGY OF TISSUE CULTURE

2.1 Selection of Plant Material (Explant)

The tissue culture process for aquascaping plants typically involves several steps, Plant explants, such as leaves, stems, or meristems, are sterilized and positioned on a nutrient-rich media in an aseptic culture, which is the initial step. An explant is the term for the plant tissue used in culture. The effectiveness of the explant is influenced by its position on the plant, as well as the plant's age or developmental stage (Katiyar & Chandel 1998). Explants that include shoot primordia, such as meristems, node buds, or shoot apices, are generally favored. Additionally, explants taken from younger or juvenile plants tend to be more successful.

Leaf explants are largely used in tissue culture as they have good potential to produce callus

and shoots in many plants. Leaf explants of *Hygrophila polysperma* were cultured on MS medium with varying concentrations of Kin or TDZ, either with or without 0.10 mg/L IBA. Direct adventitious shoot formation, occurring without intermediate callus formation, began at the leaf tips on media containing both Kin-IBA and TDZ-IBA. After one week of cultivation, distinct shoot buds appeared on the tips and edges of the leaf explants (Karatas et al, 2013a). Microsorium pteropus was established successfully from leaf explant (Suwannamali et al. 2024). *Centella asiatica* was successfully cultured using leaf explants on MS medium along with BA and NAA and also obtained 81.6% regeneration using stem node explant (Mohapatra et al. 2008).

Shoot tips are fastly dividing hence it is widely used in tissue culture. Regeneration potential of shoot tip explants of *A. heterophylla* was explored on MS medium containing 0.1 % (w/v) activated charcoal (AC) and supplemented with 6-benzylaminopurine (BAP) at different concentrations (Rittirat et al. 2021). Regeneration of *Anubias barteri var. Nana* was accomplished via organogenesis using shoot tip cultures. Multiple shoots were successfully induced from the shoot tips when grown on a modified MS medium enriched with BA and kinetin. For micro shoot induction and leaflet production, nodal explants proved to be more effective than shoot tips in three aquatic species: *Lobelia cardinalis*, *Staurogyne repens*, and *Alternanthera reineckii* (Karimi et al. 2023). Maximum number of shoots (25.33) were produced in *Hygrophila polysperma* using cultured shoot tip as explant in liquid MS medium and 21.67 shoots from first nodal segment explant (Çinar et al. 2013).

Another explant generally used in tissue culture of aquascaping plant is stem node. An ornamental aquatic plant *Ludwigiana* sp. had been successfully cultured from stem nodes containing single nodes with ± 1 cm (Lestari & Yunita 2023). In *Hedychium coronarium* rhizome bud explants cultured in BA, KIN, and TDZ, recorded maximum number of 14.21 shoots per explant on medium enriched with 1.0 mg/L TDZ and 12.89 cm shoot length on medium provided with 1.0 mg/L BA. The plants were rooted on liquid $\frac{1}{2}$ x MS medium containing 1.0 mg/L NAA with acclimatization (Verma & Bansal 2014b).

2.2 Surface Sterilization

Sterilizing explants is a crucial step for effective *in vitro* micropropagation and can be achieved

using various agents such as sodium hypochlorite, calcium hypochlorite, ethanol, mercuric chloride, hydrogen peroxide, and silver nitrate (Mihaljević et al. 2013). For aquatic plants, effective sterilization is essential before micropropagation, as entire plants or their parts are subjected to sterilization and used as explants. It is vital to ensure that explants are sterilized without significant damage, making the choice of sterilizing agent critically important. Inadequate sterilization can impede micropropagation and lead to increased contamination. The choice of sterilizing agent is influenced by the morphological characteristics of the plant parts, such as tissue hardness or softness. It is vital to ensure that explants are sterilized without significant damage, making the choice of sterilizing agent critically important. Inadequate sterilization can impede micropropagation and lead to increased contamination (Srivastava et al. 2010). For aquatic plants, hydrogen peroxide (H₂O₂) is frequently used due to its lower damage potential and effectiveness, as seen in its application to species like Water Hyssop, Dwarf Hygro, and Coontail (Karatas et al. 2013a). For water lettuce, effective sterilization has been achieved using either 1.0-3.0% diluted commercial bleach or 8-24% diluted H₂O₂. For surface sterilization of aquatic plants, such as *Bacopa monniera*, sodium hypochlorite (NaOCl) has been shown to be more successful than mercuric chloride (HgCl₂) (Tiwari et al. 1998). The performance of H₂O₂ in sterilizing *H. callitrichoides* was superior. 37 explants, or 550 individual plants (82.22%), out of 45 shoot-clump explants treated with H₂O₂ were successfully sterilized and established *in vitro* when cultivated on MS media without growth regulators (Barpete et al. 2015). The best surface sterilization for *Staurogyne repens* was obtained by utilizing 0.1% mercuric chloride for 5 minutes; this produced about 60% of the explants free of contamination. On the other hand, 1% sodium hypochlorite and 5% chloramine B were either harmful to the plants or did not work to stop microbial growth.

2.3 Preparation of Culture Media

In order to support the growth and development of aquascaping plants *in vitro*, a nutrient-rich environment must be prepared for the tissue culture of the plants. The first step in the process is to choose a suitable base medium. Two options are Gamborg's B5 medium and Murashige and Skoog (MS), both of which offer a well-balanced combination of vital nutrients. In

plant tissue culture, macronutrients such as nitrogen, potassium, calcium, magnesium, phosphorus, and sulfur make up around 80–90% of the total salt content in growth media, with nitrogen and potassium being the most prevalent. Micronutrients, including iron, manganese, zinc, boron, copper, and molybdenum, are present in much lesser quantities, often less than 1% of the overall concentration. In addition to them, though they make up a very little portion of the formulation overall, vitamins like as thiamine (B1), pyridoxine (B6), and organic compounds like sucrose are added to the medium to encourage growth and development.

The preparation involves dissolving these components in distilled water and adjusting the pH to approximately 5.8-6.0 using hydrochloric acid or sodium hydroxide. After achieving the desired pH, the medium is typically gelled with agar or Gelrite, which solidifies the medium and supports the plant tissues. The prepared medium is then sterilized by autoclaving at 121°C for 15-20 minutes to eliminate any potential contaminants. Once cooled, the sterile medium is poured into culture vessels, ready for inoculating the plant material. This carefully controlled environment is crucial for successful tissue culture, allowing aquascaping plants to grow and develop into healthy, viable specimens.

Use of liquid medium is also common in aquascaping plants culture. Liquid MS media is used in both rooting and shooting media.

2.4 Shoot Initiation

One of the most practical and economically successful techniques for propagating plants is micropropagation. The totipotency of plant cells—which enables them to regenerate entire plants in vitro—is the basis for this procedure. Since direct organogenesis is thought to be a more reliable method, it is frequently chosen for creating clonal plants that retain genetic consistency (Sandhu et al. 2018). This intricate method involves a combination of physical and chemical factors (Chand & Sahrawat 2002) and typically begins with the shoot or root meristem of the explant (Altman & Loberant 1998). The process is mainly regulated by the interaction of internal and external plant growth regulators. The genotype of the plant, the kind of explant, and its physiological state are some of the variables that affect these regulators' levels. Furthermore, the

kind and concentration of plant growth regulators employed in the culture media are important factors in determining the course and outcome of organogenesis (Amer & Omar 2019). The selection of appropriate shooting media is a critical factor in the success of micropropagation, as it can influence the growth, development, and genetic stability of the regenerated plants (Zhou & Wu 2006 and Abdalla, et al. 2022).

BAP is widely used in aquascaping plants, BAP promotes shoot multiplication and can be optimized depending on the plant species. Most aquatic plants require standard BAP concentrations of 2 to 5 mg/L; however, these can be adjusted based on the responses of individual plants. In comparison to the higher 4 mg/L BAP concentration, the medium supplemented with 1 mg/L BAP for *Cryptocoryne beckettii*, *Cryptocoryne lutea*, and *Rotala rotundifolia* induced satisfactory shoot proliferation and adventitious root formation, resulting in better root number and length. In contrast, the medium supplemented with 4 mg/L BAP produced higher rates of shoot multiplication but fewer roots in general (Micheli et al. 2006). For shoot multiplication of *Anubias barteri* var. *nana petite*, the optimal medium was MS medium supplemented with 0.2 mg/L BAP, while lowering the BAP concentration from 0.2 mg/L proved effective for promoting shoot elongation (Sheeja et al. 2015).

Another cytokinin that encourages branch growth is kinetin; however, in some aquascaping plants, it is frequently less successful than BAP. All three aquatic plants *Lobelia cardinalis*, *Staurogyne repens*, and *Alternanthera reineckii* proliferated most quickly in the MS medium containing 2 mg/L kinetin (Karimi et al. 2023). When administered with BAP or other cytokinins, kinetin's efficacy can be increased (Smith et al. 2019). *Anubias barteri* var. *Nana* was effectively regenerated by organogenesis with the use of shoot tip cultures. Cultured on a modified MS medium supplemented with BA and kinetin, several shoots emerged from the shoot tips. The ideal quantity of green shoots, averaging five shoots, was achieved using MS medium containing 3 mg/L BA. The best conditions for promoting the rooting of the regenerated shoots were MS medium supplemented with a single treatment of kinetin or without plant growth regulators. When the restored plants were put in field settings, they showed 100% acclimation and survival (Kanchanapoom et al. 2012).

Combination of auxins and cytokinins are also used in shooting in case of some plants. Using single-node shoot explants on LS media supplemented with 20 μ M BA and 0.5 μ M NAA, the maximum shoot proliferation of *Cryptocoryne lucens* was attained, with an average of 7.7 shoots per explant (Jenks et al. 2000). The best medium for *Lindernia antipoda* culture initiation and establishment was Half MS, which contained 1 mg/L of benzylaminopurine (BAP) (Rittirat et al. 2021). An alternative effective medium was 3.0 mg/L BAP, which yielded 2.40 ± 0.24 shoots per explant. The highest shoot regeneration in *Lilaeopsis brasiliensis* was observed with 1.5 mg/L NAA and 0.5mg/L BAP (Jendy et al. 2019). For shoot induction, cytokinin and auxin combinations are also utilized. The combination of 1.0 mg/L NAA and 1.0 mg/L BAP improved shoot regeneration, yielding 3.60 ± 0.24 shoots per explant and 100% of explants regenerating new shoots within 60 days. For the *Lysimachia* species, MS medium enhanced with BAP and NAA 3 was the optimal shooting medium. The best medium for *L. christinae* and *L. rubinervis* contained 3.0-5.0 mg/l BAP and 0.1 mg/l NAA, resulting in the maximum number of shoots per explant (12.25–17.20) and 100% frequency of regeneration (Zheng, et al. 2009).

2.5 Root Initiation

Root initiation is a crucial stage in the micropropagation process, which creates new plants in vitro from tiny tissue samples. Plant species conservation, genetic research, and mass propagation all make use of this technology. Rooting hormones, sometimes referred to as auxins or rooting regulators, are essential for the growth of roots from plant tissues in tissue culture. In vitro cultivated plant tissues, these hormones aid in the induction and development of roots. Since it promotes both the initiation and elongation of roots, indole-3-butyric acid (IBA) is frequently used to stimulate rooting in a variety of plant species. It is usually given at concentrations ranging from 0.1 to 5 mg/L and works especially well in woody plants and certain herbaceous species. The Murashige and Skoog (MS) nutrition medium supplemented with 0.25 mg/L Indole Acetic Acid (IAA) was shown to be the most efficient rooting medium for *Pogostemon erectus* (Dalzell) Kuntze. This effectively encouraged root formation and ensured the successful establishment of the plantlets (Dogan 2022). Massive formation of roots was observed in *Staurogyne repens* with use of a liquid MS media with addition of indole-

3-acetic acid at concentration of 0.2 mg/l (Sereda et al.2017).

Indole-3-Butyric Acid (IBA) is widely used to induce rooting in various plant species; it facilitates both root initiation and elongation, with concentrations typically ranging from 0.1 to 5 mg/L, and is particularly effective for woody plants and some herbaceous species. The best medium for establishing dwarf hygrophila (*Hygrophila polysperma*) was Murashige and Skoog (MS) medium enhanced with 0.20–1.00 mg/L of Indole-3-Butyric Acid (IBA). *Bacopa monnieri* in vitro cultivars were rooted in basal media; however, the addition of IBA greatly enhanced the rooting at 4.9 μ m levels (Tiwari, et al. 2001). The greatest quantity and longest roots of *Limnophila aromatica* were found in MS media containing 0.25 mg/ml IBA (Dogan 2019).

In many plant species, Naphthalene acetic acid (NAA) increases root production and encourages root initiation. It is frequently employed at concentrations between 0.1 and 5 mg/L and is frequently coupled with other hormones to improve rooting effectiveness. Although it can occasionally encourage the formation of roots, 2, 4-Dichlorophenoxyacetic Acid (2, 4-D) is mainly employed as a synthetic auxin for callus induction and cell elongation. It is usually applied at concentrations between 0.1 and 2 mg/L and is more frequently used for callus induction than direct rooting. *Anubias Heterophylla* had more roots when auxin, α -Naphthalene acetic acid (NAA), or 2, 4-Dichlorophenoxy acetic acid (2, 4-D) were added (Rittirat et al. 2021).

2.6 Acclimatization

The effective transition of aquascaping plants grown in vitro from a controlled tissue culture environment to natural or semi-natural settings requires acclimatization. Generally speaking, plants created by tissue culture techniques are more expensive than those cultivated using conventional approaches. It takes a lot of time and work to move plantlets from growing containers to field settings. Depending on the species or even distinct kinds, there can be differences in the complexity and requirements of this transition. Plants frequently experience structural and physiological changes throughout the adaptation phase. This procedure guarantees the plantlets' survival and growth outside of the sterile culture environment while also assisting them in acclimating to external circumstances. Start by progressively exposing the plantlets to

surroundings that are different from the sterile culture.

This can be achieved in a controlled atmosphere by gradually lowering the humidity and raising the light intensity. To reduce stress, place the plantlets in an enclosure or chamber with a high humidity level. A covered tray, a misting system, or a greenhouse with controlled humidity can all help achieve this. For a duration of one to two weeks, gradually reduce the humidity levels to facilitate the plantlets' adaptation to the reduced humidity of their surroundings. To prevent photo damage, progressively raise the light intensity, starting at lower light levels and working your way up to full light. Depending on the species, aquarium plants will require anywhere between 300 to 6,000 lux. Make sure you progressively adjust the temperature to the parameters of the intended growing environment. The most commonly used tool to control the temperature in an aquarium is a submersible aquarium heater with an integrated thermostat. This heater is placed inside the tank, and the thermostat allows for precise temperature regulation, ensuring the water stays within the desired range for the fish and other aquatic life.

In the micropropagation process, hardening media is essential, especially for acclimating in vitro produced plantlets to ex vitro environments. Bridging the gap between the regulated in vitro culture and the natural or semi-natural circumstances in which the plants will ultimately be grown is the main objective of hardening media. Plants used in aquascaping are frequently produced in nutrient-rich, controlled environments, so reintroducing them to natural or semi-natural settings can be challenging. In order to help these aquatic or semi-aquatic species adjust from in vitro circumstances to aquarium or terrestrial settings, hardening media for aquascaping plants must satisfy their specific needs. Various hardening medium are used to help aquascaping plants that are grown in vitro acclimate to aquarium or terrestrial habitats more easily. Plant growth is supported by aqua soil, which also supplies vital nutrients. Good drainage and stability are guaranteed by gravel; well-liked varieties include Seachem Flourite Gravel and CaribSea Eco-Complete. With items like Seachem Flourite Sand and Pool Filter Sand, sand provides a natural appearance while supporting fragile roots. Nutrient retention is improved by clay-based substrates like ADA Aqua Soil and Fluval Plant and Shrimp Stratum. Perlite and vermiculite enhance aeration and

moisture retention, whereas peat moss contributes moisture and acidity. Excellent drainage is provided by hydroton, and a well-balanced structure is guaranteed by soil-based mixes. Customized growth conditions can be established to fit the needs of individual plants by combining different substrates. *Nymphoides indica* was effectively acclimated by using clay pots with a 3:1 mixture of peat moss and decomposed cow manure. These pots were then submerged in culture tanks 10 cm below the water's surface and exposed to full sunlight (Jenks et al. 2000). For *Cunila galioides*, rooted plantlets were acclimated by moving them into plastic chambers containing a sterilized sand-soil mixture (1:1). Initially covered with a plastic cap, the cap was gradually removed over a period of two weeks before the plants were transitioned to a greenhouse and later to outdoor conditions (Fracaro et al. 2001). Hardening of *Bacopa monnieri* plantlets is done by placing them in polybags with a 1:1:1 mixture of sand, farmyard manure, and soil, irrigating them with half-strength MS medium in a mist chamber for three weeks, then transitioning to an open shade house with tap water irrigation for one month before field transfer (Sharma et al. 2012). *Ludwigiana repens* plantlets were effectively adapted in aquarium tanks with a 100% success rate in a study by Öztürk et al. (2004). In fiber tanks housed within a net house, *Bacopa caroliniana*, *Anubias minima*, *Aponogeton ulvaceus*, *Rotala rotundifolia*, and *Nymphoides cristatata* were successfully hardened utilizing a substrate mixture of pond soil and coir fibers (Navya 2013). A soil mix similar to natural conditions (peat:clay in a 1:1:10 ratio) produced superior shoot and root growth for acclimating *Cryptocoryne beckettii*, *Cryptocoryne lutea*, and *Rotala rotundifolia* in aquarium tanks than the commercial Compo Cactea© mix (Abdalla et al. 2022).

3. CHALLENGES IN MICROPROPAGATION OF AQUASCAPING PLANTS

Aquascaping plant micropropagation has a number of difficulties, such as the requirement for specialized nutrient formulations to promote healthy plant development and the necessity for exact control over growth conditions. Plant responses to growth regulators and medium compositions are inherently variable, which can make successful plant regeneration more difficult. Additionally, because there is a significant chance of contamination from

microbial sources, preserving sterility throughout the culture phase is essential yet challenging. Acclimatization problems are frequently the consequence of moving from in vitro to ex vitro settings; plants may find it difficult to adjust to their new surroundings, which can result in high mortality rates. Additionally, it can be expensive and hard to optimize techniques for various plant species, necessitating a great deal of trial and error. To overcome these obstacles, careful attention to detail, thorough testing, and adaptation of protocols to the specific needs of each aquascaping plant species.

Micropropagation has become a crucial technique for the efficient and sustainable cultivation of ornamental aquatic plants, offering significant benefits such as the production of numerous uniform, disease-free specimens and the capacity for rapid multiplication. However, this method also presents several challenges, including the necessity for precise control over growth conditions, the risk of contamination, and the complexity of tailoring protocols to different species. Effective micropropagation demands a thorough understanding of plant physiology along with a customized approach to growth media, regulators, and acclimatization strategies. Ongoing advancements in this field are improving the efficiency and success rates of plant propagation, which supports more sustainable practices in aquascaping and helps conserve valuable aquatic species. Future research should aim to address existing challenges, refine protocols, and explore new techniques to further enhance the micropropagation of aquascaping plants.

4. CONCLUSION

Micropropagation has emerged as a key method for the productive and sustainable development of ornamental aquatic plants. It has a number of advantages, including the ability to multiply quickly and produce a large number of uniform, disease-free specimens. Nevertheless, there are a number of drawbacks to this approach as well, such as the need for exact control over growing conditions, the possibility of contamination, and the difficulty of customizing protocols for various species. Effective micropropagation demands a thorough understanding of plant physiology along with a customized approach to growth media, regulators, and acclimatization strategies. Ongoing advancements in this field are improving the efficiency and success rates of plant propagation, which supports more

sustainable practices in aquascaping and helps conserve valuable aquatic species. Future research should aim to address existing challenges, refine protocols, and explore new techniques to further enhance the micropropagation of aquascaping plants.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

I hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdalla, N., El-Ramady, H., Seliem, M. K., El-Mahrouk, M. E., Taha, N., Bayoumi, Y., ... & Dobránszki, J. (2022). An academic and technical overview on plant micropropagation challenges. *Horticulturae*, 8(8), 677.
- Altman A, Loberant B. (1998). Micropropagation: clonal plant propagation in vitro. In Altman A, ed. *Agricultural biotechnology*. New York: Marcel Dekker. 19–42
- Amer A and Omar H. (2019). In-vitro propagation of the multipurpose Egyptian medicinal plant *Pimpinella anisum*. *Egypt. Pharm. J.* 18:254–262. DOI: 10.4103/epj.epj_12_19.
- Baby, G., Varma, P., Patnaik, K. and Rahman, A. M. (2024). Aquascaping: Where Nature Meets Art in Your Home Aquarium. *Farm Chronicle*, 03(09), 36-40.
- Barpete, S., Özcan, S. F., Aasim, M., & Özcan, S. (2015). In vitro high frequency regeneration through apical shoot proliferation of *Hemianthus callitrichoides* Cuba'-a multipurpose ornamental aquatic plant. *Turkish Journal of Biology*, 39(3), 493-500.
- Chand, S. and Sahrawat, A.K. (2002). Somatic embryogenesis and plant regeneration from root segments of *Psoralea corylifolia* L., an endangered medicinally important plant. *In Vitro Cellular & Developmental Biology-Plant*, 38(1):33-38.
- Çınar A, Karataş M, Aasim M (2013) High frequency plant regeneration of dwarf hygro (*Hygrophila polysperma* [Roxb.] T.

- Anderson) on liquid culture. *J Appl Biol Sci* 7:75–78.
- Dogan, M. (2019, December). Effect of Different Indole-3-Butyric Acid and Naphthalene Aetic Acid on in vitro Rooting of *Limnophila aromatica* (Lamk.) Merr. In *CONGRESS PROCEEDINGS SERIES* (p. 187).
- Dogan, M. (2022). In vitro micropropagation of *Pogostemon erectus* (Dalzell) Kuntze in liquid culture medium. *Natural and Engineering Sciences*, 7(1), 80-88
- Espinosa-Leal, C. A., Puente-Garza, C. A., & García-Lara, S. (2018). In vitro plant tissue culture: means for production of biological active compounds. *Planta*, 248, 1-18.
- Fracaro, F., & Echeverrigaray, S. (2001). Micropropagation of *Cunila galioides*, a popular medicinal plant of south Brazil. *Plant Cell, Tissue and Organ Culture*, 64, 1-4.
- Hussain, A., Qarshi, I. A., Nazir, H., & Ullah, I. (2012). Plant tissue culture: current status and opportunities. *Recent advances in plant in vitro culture*, 6(10), 1-28.
- Jabir, T., George, S., Raj, A., Lakshmi, S., & Joseph, A. (2016). Micropropagation and in vitro flowering of an ornamental aquarium plant *Lindernia antipoda* (L.) Alston. *Intl. J. Aquaculture*, 6(8), 1-10.
- Jendy, J. A., Ha, H. C., Zainuddin, R., Zawawi, D. D., & Ing, N. S. (2019). Micropropagation of aquatic plant Brazilian micro sword (*Lileaopsis brasiliensis*). *Journal of Agrobiotechnology*, 10(1S), 29-34.
- Jenks, M. A., Kane, M. E., & McConnell, D. B. (2000). Shoot organogenesis from petiole explants in the aquatic plant *Nymphoides indica*. *Plant Cell, Tissue and Organ Culture*, 63, 1-8.
- Jenks, M. A., Kane, M. E., & McConnell, D. B. (2000). Shoot organogenesis from petiole explants in the aquatic plant *Nymphoides indica*. *Plant Cell, Tissue and Organ Culture*, 63, 1-8.
- Kanchanapoom, K., Chunui.P., and Kanchanapoom, K. 2012. Micropropagation of *Anubias barteri* var. Nana from shoot tip culture and the analysis of ploidy stability. *Not. Bot. Hort. Agrobo.* 40: 148-151.
- Karatas M, Aasim M, Dogan M, Khawar KM. 2013a. Adventitious shoot regeneration of the medicinal aquatic plant water hyssop (*Bacopa monnieri* L. Pennell) using different internodes. *Archives of Biological Sciences, Belgrade.* 65:297-303.
- Karimi Alavijeh, M., Safi, S., & Zarei, A. (2023). An efficient method for economic micropropagation of three aquatic plant species (*Lobelia cardinalis*, *Staurogyne repens*, and *Alternanthera reineckii*). *Aquaculture International*, 31(3), 1623-1636.
- Karimi Alavijeh, M., Safi, S., & Zarei, A. (2023). An efficient method for economic micropropagation of three aquatic plant species (*Lobelia cardinalis*, *Staurogyne repens*, and *Alternanthera reineckii*). *Aquaculture International*, 31(3), 1623-1636.
- Katiyar SK, Chandel G (1998) High frequency somatic embryogenesis from immature inflorescence of *Coix aquatica* Roxb. *Plant Tissue Cult* 8:131–137
- Kunene, E. N., Oseni, T. O., Wahome, P. K., Masarirambi, M. T., McCubbin, M. J., Dlamini, P. S., ... & Zwane, M. G. (2018). Effects of plant growth regulators and explant type on the in vitro micropropagation of wild ginger (*Siphonochilus aethiopicus* (Schweif.) BL Burt.). *Adv Med Res*, 6, 54-63.
- Lestari, E. G., & Yunita, R. (2023). The Formula media in vitro Propagation and Conservation of *Ludwigia* sp. *Journal of Tropical Biodiversity and Biotechnology*, 8(1), 75947.
- Martin, M. (2013). *Aquascaping: Aquarium landscaping like a pro*. Ubiquitous Publishing.
- Micheli, M., De Gasperis, A., Prospero, F., & Standardi, A. (2006). Micropropagation of three species of aquatic plants. *Agricoltura Mediterranea*, 136(1), 46.
- Mihaljević, I., Dugalić, K., Tomaš, V., Viljevac, M., Pranjić, A., Čmelik, Z., & Jurković, Z. (2013). In vitro sterilization procedures for micropropagation of 'Oblačinska'sour cherry. *Journal of Agricultural Sciences, Belgrade*, 58(2), 117-126.jlj
- Mohapatra H, Barik DP, Rath SP (2008) In vitro regeneration of medicinal plant *Centella asiatica*. *Biol Plant* 52:339.
- Navya, R. (2013). *Integrated culture of fish with micropropagated plants in a recirculatory system* (Doctoral dissertation, KUFOS).
- Öztürk, M., Khawar, K. M., Atar, H. H., Sancak, C., & Ozcan, S. (2004). In vitro micropropagation of the aquarium plant *Ludwigia repens*. *Asia Pacific Journal of Molecular Biology & Biotechnology*, 12(1 & 2), 21-25.

- Pierik R.L.M., Ruibing M.A. (1997) - Developments in the micropropagation industry in The Netherlands. *Plant Tissue Culture and Biotechnology* Vol. 3.
- Rittirat, S., Thammasiri, K., & Klaocheed, S. (2021). In vitro rapid multiplication of a highly valuable ornamental aquatic plant *Anubias heterophylla*. *Trends in Sciences*, 18(19), 3-3.
- Sandhu M, Wani SH, Jiménez VM. (2018). In vitro propagation of bamboo species through axillary shoot proliferation: a review. *Plant Cell Tiss Organ Cult*. 132:27–53.
- Sereda, M. M., Lutsenko, E. V., Chokheli, V. A., Vereschagina, A. V., Rachkovskaya, K. Y., Lysenko, V. S., & Varduny, T. V. (2017). A method for microclonal propagation of *Staurogyne repens* in tissue culture. *Journal of Plant Sciences*, 12(1), 17-21.
- Sharma, N., Satsangi, R., Pandey, R., Singh, R., Kaushik, N., & Tyagi, R. K. (2012). In vitro conservation of *Bacopa monnieri* (L.) using mineral oil. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 111, 291-301.
- Sheeja, G. E., Joseph, A., & Korath, A. (2015). In vitro propagation of an ornamental aquatic plant, *Anubias barterii* Var. Nana petite. *Int. J. Curr. Sci*, 18, 1-12.
- Smith, S. C., Warren, L. M., & Cooley, L. D. (2019). Maintaining a methods database to optimize solid tumor tissue culture: Review of a 15-year database from a single institution. *Cancer Genetics*, 233, 96-101
- Srivastava N, Kamal B, Sharma V, Negi YK, Dobriyal AK, Gupta S, Jadon VK. 2010. Standardization of sterilization protocol for micropropagation of *Aconitum heterophyllum*- an endangered medicinal herb. *Academic Arena*. 2:37-42.
- Suwannamali, W., Wang, K. T., Su, C. C., Kantha, P., Tzean, Y., & Wu, T. M. (2024). Optimizing Green Globular Body Induction for Micropropagation of *Microsorium pteropus* 'Windeløv'. *Horticulturae*, 10(7), 673.
- Tiwari, V., Deo Singh, B., & Nath Tiwari, K. (1998). Shoot regeneration and somatic embryogenesis from different explants of Brahmi [*Bacopa monnieri* (L.) Wettst.]. *Plant Cell Reports*, 17, 538-543.
- Tiwari, V., Tiwari, K. N., & Singh, B. D. (2001). Comparative studies of cytokinins on in vitro propagation of *Bacopa monnieri*. *Plant cell, tissue and organ culture*, 66, 9-16.
- Verma M, Bansal YK (2014b) Induction of somatic embryogenesis in endangered butterfly ginger *Hedychium coronarium* J. Koenig. *Indian J Exp Biol* 50:904–909
- Zheng, W., Xu, X. D., Dai, H., & Chen, L. Q. (2009). Direct regeneration of plants derived from in vitro cultured shoot tips and leaves of three *Lysimachia* species. *Scientia Horticulturae*, 122(1), 138-141.
- Zhou, L. G., & Wu, J. Y. (2006). Development and application of medicinal plant tissue cultures for production of drugs and herbal medicinals in China. *Natural Product Reports*, 23(5), 789-810.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125306>