



***In vitro* Study on Anti-salmonella Activities of *Boerhaavia diffusa* (L. syn) Leaf Extract**

Ologun, Oluwabusayo^{1*}, Dada, Ebenezer Oluyemi¹ and Ajayi, Kehinde Oluyemi¹

¹*Department of Microbiology, Federal University of Technology, Akure, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author OO designed the study, wrote the protocol and first draft of the manuscript. Author DEO also designed the study and managed the literature search. Author AKO performed the statistical analysis, interpreted the data and managed the literature search. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPR/2019/v3i130081

Editor(s):

- (1) Dr. Jasini A. Musa, Department of Veterinary Microbiology, University of Maiduguri, Nigeria.
(2) Dr. Nagesh Peddada, Department of Biophysics, University of Texas Southwestern Medical Center, USA.

Reviewers:

- (1) R. Prabha, Dairy Science College, India
(2) Bogumil E. Brycki, Adam Mickiewicz University, Poland.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49728>

Original Research Article

Received 20 April 2019

Accepted 29 June 2019

Published 22 July 2019

ABSTRACT

Various strategies have been employed in the treatment and management of Salmonella infection however, Salmonella strains have gained resistance to antibiotics. This study was to determine in vitro anti-Salmonella activity of *Boerhaavia diffusa* leaf extract against clinical isolate of *Salmonella typhi* and *Salmonella typhi* ATCC 14028. The aqueous and ethanol extracts of *B. diffusa* were studied for their antibacterial activity against pathogenic *Salmonella typhi*. This study was carried out between April and September 2018. The in vitro antibacterial activity was performed by agar well diffusion method and broth dilution using spectrophotometric method and the results were expressed as the average diameter of zone of inhibition of bacterial growth around the well and optical density respectively. It was observed that aqueous extract exerted slightly higher activity than ethanolic extract as revealed by the mean diameter of zone of inhibitions at a concentration of 200 mg/ml, the aqueous extract had 35.21±0.47 mm (*Salmonella typhi* ATCC 14028) compared with ethanol extract 26.41±0.32 mm (clinical). However, in the broth dilution method, ethanol extract significantly (p=0.05) reduced the cell, at 48 hours, the optical density of clinical isolate of *S. typhi* treated at concentration of 200 mg/ml of extract was 0.47±0.02 nm while at the same concentration of extract, aqueous extract had an optical density of 0.52±0.11 nm respectively. The phytochemical assay revealed that tannin (5.18±0.02 mg/g) and quinone (8.45±0.13 mg/g) in

*Corresponding author: Email: ologunoluwabusayo@gmail.com;

ethanol extract was significantly ($p=0.05$) higher than aqueous extract while saponin (14.18 ± 0.06 mg/g) was higher in the aqueous extract. The ethanol and aqueous extracts of leaves of *B. diffusa* whole plant exhibited significant antibacterial activity against both clinical and typed *Salmonella typhi*. Therefore, the plant extract could be used for the treatment of Salmonellosis, however, the in vivo studies is needed to ascertain the safety of the extract.

Keywords: Anti-salmonella activity; plant extracts; agar well diffusion; broth dilution; salmonella strains.

1. INTRODUCTION

The bacterium *Salmonella typhi* causes typhoid fever [1,2]. The bacterium is a gram-negative, motile, non-sporing, non-capsulated bacillus that can be contracted through contaminated water, milk, food or fruits and vegetables or via convalescent or chronic carriers [3]. It has also been linked with zoonotic transmission via reptiles and common domestic pets [4]. *Salmonella enterica*, which is a group of Gram-negative bacterial pathogens capable of infecting humans and animals, cause significant morbidity and mortality worldwide [5]. Certain serotypes adapted to humans, such as *Salmonella typhi* (*S. typhi*) and *Salmonella paratyphi* (*S. paratyphi*), usually cause severe diseases in humans, such as enteric fevers (typhoid and paratyphoid fevers). In most endemic areas like Africa, Asia, and Latin America, approximately 90% of enteric fever is typhoid. This disease is an important global health problem with an estimated 16 million cases and 600 000 deaths each year.

Various strategies have been employed in the treatment and management of *Salmonella* infection. Fluoroquinolones and tetracyclines are most commonly used to treat *Salmonella* infections. However, *Salmonella* strains resistant to these antibiotics have been reported in Korea and other countries [6]. One major concern to public health has been the global dissemination of *S. typhimurium* Definitive Type 104, which is resistant to cotrimoxazole, nalidixic acid and ampicillin [7]. The rise in antibiotic-resistant strains has led to increased interest in the use of plant materials to develop new effective drugs [6]. Moreover, conventional antityphoid drugs are becoming more and more unavailable to the common man in Africa due to increased cost [8].

The rise in antibiotic-resistant strains has led to increased interest in the use of plant materials to develop new effective drugs [6]. It has been reported that 80% of the world population are rural dwellers and rely on medicinal plants for their daily medications, also, plants have been reported to have minimal or no side effects

compared to antibiotics [9,10]. *Boerhaavia diffusa* (Spreading Hogweed in English), belonging to the family of the Nyctaginaceae, is mainly a diffused perennial herbaceous creeping weed of India (known also under its traditional name as *Punarnava*). *Boerhaavia diffusa* is traditionally known in Nigeria as *Etiponla* in Yoruba, *Azeigwe* in Igbo and *Babba-juju* in Hausa. *B. diffusa* is a perennial creeping weed, prostrate or ascending herb, up to 1 m long or more, having spreading branches [11].

The root, leaves, aerial parts and the whole plant of *B. diffusa* (L. syn) are used worldwide for the treatment of a number of disorders e.g. liver complaints, kidney disorders, rheumatism e.t.c. [12]. The quest to identify and isolate novel phyto-compounds from *B. diffusa* has led many researchers to discover various compounds such as flavonoids, alkaloids, glycosides, steroids, triterpenoids, lipids, lignans, carbohydrates, proteins, and glycoproteins from its leaves, stems, seeds and roots [13]. Sourav [14], explored the Antibacterials from *Boerhaavia diffusa*. In his study, the chloroform and alcohol extracts of the plant were screened against six bacteria viz *Staphylococcus aureus*, *Escherichia coli*, *Proteus mirabilis*, *Salmonella typhimurium*, *Pseudomonas aeruginosa* and *Klebsiella aerogenosa*. Chloroform extract showed activity against *E. coli*, *S. typhimurium* and *P. aeruginosa* while the alcohol extract was active against *P. mirabilis* and *S. typhimurium*. The present study was undertaken to further investigate the antibacterial activity of *Boerhaavia diffusa* on typed and clinical strains of *Salmonella typhi* with the view to provide scientific evidence for its application as a medicinal plant.

2. MATERIALS AND METHODS

2.1 Collection of Leaves of *Boerhaavia*

Fresh leaves of *Boerhaavia diffusa* (Plate 1) were collected from the School of Health Technology, Oda Road, Akure, and identified in

the Department of Crop, Soil and Pest Management, Federal University of Technology, Akure Ondo State.



Plate 1. Photograph of *Boerhaavia diffusa* leaf

2.2 Preparation of Plant Extract

2.2.1 Aqueous extraction

The aqueous extractions of the water-soluble ingredient were carried out using the filter method. A 2 g of each of the grounded leaves were extracted by successive soaking for 2 days using 50 ml of distilled water in a 250 ml sterile conical flask. The extracts were concentrated in vacuum at 60°C and stored in universal bottles and refrigerated at 4°C prior to use [15].

2.2.2 Ethanol extraction

The organic solvent leaf extract was prepared by 2 g of plant mixture with ethanol and kept for two days. The extract was concentrated to one-fifth volume, filter sterilized and stored at 4°C [15].

2.3 Test Organism

The clinical bacterial strains were obtained from the Department of Microbiology, Federal University of Technology Akure. Clinical *Salmonella typhi* and typed (ATCC 14028) *Salmonella typhi* were used. The isolates were confirmed based on cultural, morphological and biochemical characteristics following standard methods of identifying *Salmonella typhi* [16]. The bacterial strain was grown in nutrient broth for 12-18 hours at 37°C on a rotary shaker. Cells were grown at 37°C for 18 hours and cultures were kept at 4°C.

2.3.1 Antimicrobial susceptibility tests

Standardization of the inoculums: The inoculum was prepared by inoculating colonies of fresh test cultures into sterile distilled water. The turbidity was compared to 0.5 McFarland standard prepared according to the method of Cheesbrough [16]

2.4 Antibacterial Susceptibility Assay

The extracts were dissolved and diluted using 50% v/v dimethylsulphoxide (DMSO) to obtain different concentrations (50, 100 and 200 mg) in 1 mL. The 50 mg/ml, 100 mg/ml and 200 mg/ml of the extracts of *B. diffusa* leaves were introduced into the wells of Muller Hinton agar plate. The plates were incubated aerobically at 37°C and examined after 24 hours. The plates were examined for microbial growth inhibition and the Inhibition Zone Diameter (IZD) was measured to the nearest millimetre and compared with those produced by the commercial antibiotic ciprofloxacin which was used as control. Effect of extract on anti-*Salmonella* efficacy of the extract in the broth was also assayed using the spectrophotometric method, the absorbance of the tube was read at 620 nm [16,17].

2.5 Antibiotics sensitivity Test Using Commercial

Antibiotics sensitivity test of the bacterial isolates were determined by the disc diffusion method as described by Cheesbrough [16]. Standard inoculum of 18 hours broth was spread on Muller Hinton agar using sterile swab in triplicate. The antibiotic discs were placed on the plate at equidistance. The plates were then incubated for 24 hours at 37°C and the diameter of zone of inhibition was measured and recorded. The commercial antibiotics discs (Fondoz Laboratories Ltd, Nigeria) used were; Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg, Ciprofloxacin (CPX) 10 µg, Amoxicillin (AM) 25 µg, Augmentin (AU) 30 µg, Gentamycin (CN) 10 µg, Pefloxacin (PEF) 5 µg, Ofloxacin (OFX) 5 µg, Streptomycin (S) 10 µg and Septra (SXT) 30 µg.

Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of *Boerhaavia diffusa* Extracts: The Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the extracts were determined using the broth (tube) dilution

technique [18]. Dilutions of the extract in Mueller Hinton broth were prepared in tubes. The concentration of inoculum was also standardized to 0.5 McFarland's turbidity, The Mueller Hinton broth in tubes containing the different concentration of plant extract, 50, 100, and 200 mg/ml was then inoculated with 0.5 ml of the standardized culture. The tubes were then incubated at 37°C for 24 hours. MIC and MBC values were recorded.

2.6 Screening of Phytochemical Compounds

The various solvent extracts of the powder of leaves of *Boerhaavia diffusa* were subjected to phytochemical tests for the identification of various action constituents using the method of Marcelin et al. [17]. The following major pharmaceutical valuable phytochemical compounds were analyzed qualitatively and quantitatively; alkaloids, phenols, tannins, flavonoids, quinones, saponins, terpenoids, sterols and cardiac glycosides.

2.7 Statistical Analysis of Data

Data obtained were subjected to analysis of variance and means were compared using Duncan's New Multiple Range Test (DNMRT) with the aid of SPSS software at $p \leq 0.05$ level of significance.

3. RESULTS AND DISCUSSION

Salmonellosis and enteric fever are always a public health concern in most developing countries, which are mostly low or middle-income countries with inadequate sanitation and hygiene, particularly, regarding food, water and disposal of human excreta [17]. Different plants and their parts (flowers, buds, leaves, stem, bark, fruits, skin, pulp and root) have been used for thousands of years to enhance the flavour and aroma of food. In addition, plants are rich in a wide variety of second metabolites such as Alkaloids, Flavonoids, Phenols, which were found *in vitro* to have antimicrobial properties [17,19].

In this study, extracts of *Boerhaavia diffusa* leaves were investigated for antibacterial activity against *Salmonella typhi*. Plant extracts were used to investigate antibacterial activity against two bacterial strains (Clinical *Salmonella typhi* and *Salmonella typhi* ATCC 14028). In this study, the antibacterial activity of *B. diffusa* leaf extracts

was compared against the test bacteria with activities of model antibiotics.

The test organisms used for this study were identified based on biochemical characteristics common to *Salmonella typhi*. The result is presented in Table 1. The antibiotic sensitivity patterns of commercial antibiotics on the two strains of *S. typhi* are presented in Fig. 1. The result revealed that the zones of inhibition of antibiotics against typed isolates were higher than that of clinical isolates however, chloramphenicol had the highest inhibition against the isolates (STC=24.30±0.42 mm, STT=24.36±0.07 mm). The higher antibacterial activity of model antibiotics is not surprising since the antibiotics are in a refined state. The standard antibiotics (ampicillin, amoxicillin, ciprofloxacin, ofloxacin, chloramphenicol) used in this study are the first line drugs employed in the treatment of typhoid fever [1].

Table 1. Biochemical characteristics of *Salmonella* strains

Biochemical characteristics	<i>Salmonella typhi</i> (Clinical isolate)	<i>Salmonella typhi</i> (ATCC 14028)
Gram reaction	-ve	-ve
Shape	Rod	Rod
Motility	+ve	+ve
Catalase	+ve	+ve
Coagulase	-ve	-ve
Citrate	+ve	+ve
H ₂ S	+ve	+ve
Lactose	-ve	-ve
Glucose	+ve	+ve
Fructose	+ve	+ve
Sucrose	-ve	-ve
Galactose	+ve	+ve
Indole	-ve	-ve
Methyl red	+ve	+ve
Voges-Proskauer	-ve	-ve
Oxidase	-ve	-ve

Key: -ve= negative; +ve= positive

The results of antibacterial activity of both water and ethanol crude extracts of *B. diffusa* showed anti*Salmonella* activity on the two strains of *S. typhi* tested at different concentrations, with aqueous extract exerting slightly higher activity than ethanolic extract as revealed by mean diameter of zone of inhibitions, 200 mg/ml of aqueous extract had the highest (35.21±0.47) zone of inhibition (Fig. 2). Minimum inhibitory concentration (MIC) and minimum bactericidal concentrations (MBC) of the extracts is shown in Tables 2. The ethanol and aqueous extract had the same MIC (100 mg/ml) on the typed isolate, also, there was no difference in the MIC and

MBC of ethanol and aqueous extract on typed isolate. The aqueous and ethanol extracts exhibited different zones of inhibition against the isolates, however, the aqueous extract had higher zones of inhibition than ethanol extract. Antimicrobial action may be due to the synergistic action of different chemical constituents, some of which probably are lost upon extraction with solvent [15,17,20]. Water could be a better extraction solvent than ethanol for *B. diffusa* leaf, also, the demonstration of higher activity by the aqueous solvent may be an indication that the phytoconstituents in the plant leaves are more soluble in water than the organic solvent [21]. The antimicrobial potential of *B. diffusa* and other plants sourced from traditional healers through an ethnobotanical survey of anti-infective plants in Egbado South in Ogun State, Nigeria was previously reported by Abo and Ashidi [22]. This study also corroborates the findings of Madani and Jain [23] who reported higher anti-*Salmonella* activity in aqueous extract of *Terminalia bellerica* than chloroform and acetone extracts. It has been reported that different phytoconstituents have different degrees of solubility in different types of solvents depending on their polarity. In a traditional setting, water is the solvent largely used to prepare these concoctions.

Table 2. Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of *Boerhaavia diffusa* extracts on *Salmonella*

<i>Boerhaavia diffusa</i> extracts	Ethanol extract		Aqueous extract	
	S1	S2	S1	S2
MIC (mg/ml)	100	100	50	100
MBC (mg/ml)	50	100	50	100

Key: S1 - *S. typhi* (Clinical isolate), S2 - *S. typhi* (ATCC 14028)

The anti-*Salmonella* efficacy of *Boerhaavia diffusa* extracts in the broth was assayed and was shown in Figs. 3, 4, 5 and 6. The result presented in Fig. 3 revealed the effect of ethanol extract on clinical isolate of *S. typhi*, it was noted that the extract significantly ($p < 0.05$) reduced the cell, at 48 hours, the optical density of clinical isolate of *S. typhi* treated with 50, 100, 200 mg/ml of extract were 0.52 ± 0.03 , 0.50 ± 0.10 , 0.47 ± 0.02 nm respectively while at the same concentration of extract, aqueous extract had optical density of 0.64 ± 0.21 , 0.54 ± 0.03 , 0.52 ± 0.11 nm respectively (Fig.4). Also, the anti-*Salmonella* efficacy of *B. diffusa* ethanol extracts on a typed isolate of *S. typhi* is shown in Fig.5. It was observed that the extract significantly ($p < 0.05$) reduced the cell,




Fig. 1. Antibiotic sensitivity pattern of commercial antibiotic discs on *S. typhi* strains
 Key: STC – *Salmonella typhi* (clinical isolate), STT – *Salmonella typhi* (typed isolate), Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg, Ciprofloxacin (CPX) 10µg, Amoxicillin (AM) 25 µg, Augmentin (AU) 30 µg, Gentamycin (CN) 10 µg, Pefloxacin (PEF) 5 µg, Ofloxacin (OFX) 5µg, Streptomycin (S) 10 µg and Septra (SXT) 30 µg




Fig. 2. AntiSalmonella activity of Boerhaavia diffusa extracts

Key: AQ = Aqueous extracts of Boerhaavia diffusa, ET= Ethanolic extracts of Boerhaavia diffusa, Cpx= Ciprofloxacin




Fig. 3. Effect of Boerhaavia diffusa ethanol extract on clinical isolate

at 48 hours, the optical density was of typed isolate of *S. typhi* treated with 50, 100, 200 mg/ml of extract were 0.49 ± 0.00 , 0.48 ± 0.01 and 0.37 ± 0.12 while at the same concentration, aqueous extract had an optical density of 0.62 ± 0.03 , 0.53 ± 0.11 and 0.49 ± 0.21 nm respectively (Fig. 6). It was noted from this study that plant extracts tested by microdilution technique and the optical density was measured

after 48 hours showed that ethanol extract had higher anti-*Salmonella* activity compared to aqueous extract which was higher in values obtained from agar well diffusion technique. It could be that the bioactive components in ethanol extract did not diffuse into agar in agar well but was able to inhibit microbial cells directly in broth. This was previously reported by other findings that the active components of the extract

do not diffuse into Muller Hinton agar, however, they were able to cause inhibition of microbial cells in broth microdilution [6,24].

Both plant extracts (ethanolic and aqueous) were subjected to preliminary qualitative phytochemical evaluation. The phytochemical profiles of the two solvent extracts from the plant used in this study are presented in Table 2. The analysis revealed the presence of alkaloids, phenol, glycosides, steroids, carboxylic acid, reducing sugar, flavonoids, saponins, tannins, proteins, triterpenoids, quinines, carbohydrates and sterols. Also, tannin (5.18 ± 0.02 mg/g) and quinone (8.45 ± 0.13 mg/g) in ethanol extract were significantly ($p < 0.05$) higher than aqueous extract while saponin (14.18 ± 0.06 mg/g) was higher in the aqueous extract. The preliminary qualitative phytochemical screening carried out showed that the leaf extracts of *B. diffusa* contain vital secondary metabolites such as alkaloids, saponins, tannins and glycosides. The bioactive compounds in medicinal plants have been reported to be the active principles responsible for the pharmacological potentials of medicinal plants [25]. The presence of these chemicals in the leaves and root of these plants justify the local uses of these plants for the treatment of various ill conditions. Phytoconstituents such as saponins, phenolic compounds and glycosides have been reported

to inhibit bacterial growth and to be protective to plants against bacterial and fungal infections [6,26]. Ethanol extract of *Boerhavia diffusa* leaves possess some phytochemicals like Alkaloids, Anthraquinone, Glycoside, Flavanoids and Tannins. Saponins are natural glycosides that act as hypoglycemic, antifungal and serum cholesterol lowering agents in animals [27]. Saponins are essential elements in ensuring hormonal balance and synthesis of sex hormones [28]. Tannins are bitter polyphenolic compounds that hasten the healing of wounds. They also possess anti-diuretic and anti-diarrhoea properties [28]. Terpenoids were present in both ethanolic extract of and aqueous extracts of AOU and AFU. Terpenoids have been found to be useful in the prevention and therapy of several diseases, including cancer. Terpenoids are also known to possess antimicrobial, antifungal, antiparasitic, antiviral, anti-allergenic, antispasmodic, antihyperglycemic, anti-inflammatory and immunomodulatory properties [29]. The presence of these compounds promises its potential application in the treatment of microbial ailment. However, tannins were present in aqueous extract of but not in the ethanolic extract. Saponin and flavanoid are higher in the aqueous extract of the leaf (14.18 and 11.26 mg/g) than the ethanolic extract (6.36 and 9.98 mg/g).




Fig. 4. Effect of boerhaavia diffusa aqueous extract on clinical isolate

Table 3. Qualitative analysis of phytochemicals in *Boerhavia diffusa* leaf extracts

Phytochemical	Ethanolic extract	Aqueous extract
Alkaloids	-	+
Tannins	-	+
Flavonoids	+	+
Quinones	+	-
Saponins	+	+
Terpenoids	+	-
Sterols	+	-
Cardiac Glycosides	-	+
Phenols	+	+

Key: + = Present, - = Absent




Fig. 5. Effect of *Boerhaavia diffusa* ethanol extract on typed isolate




Fig. 6. Effect of *Boerhaavia diffusa* aqueous extract on typed isolate

Table 4. Quantitative phytochemical screening of aqueous and ethanol extracts of *B. diffusa*

Phytochemicals	Ethanolic extract	Aqueous extract
Tannins (mg/g)	5.18±0.02 ^a	3.90±0.22 ^a
Quinones (mg/g)	8.45±0.13 ^b	6.60±0.31 ^a
Saponins (mg/g)	6.36±0.24 ^a	14.18±0.06 ^b
Triterpenoids (mg/g)	8.56±0.08 ^a	8.89±0.31 ^a
Steroid (mg/g)	9.03±0.11 ^a	6.73±0.14 ^a
Glycosides (mg/g)	30.39±0.06 ^b	28.29± 0.03 ^a
Flavonoids (mg/g)	9.98±0.61 ^a	11.26±0.33 ^a

4. CONCLUSION

Most of the antibiotics used nowadays have lost their effectiveness due to the development of resistant genes in microbes. The antibiotics are sometimes associated with side effects such as hypersensitivity, immune suppression and allergic reaction.

More interest is being shown in developing alternative antimicrobial drugs for the treatment of infectious diseases without side effects. The results of our present study demonstrates anti-*Salmonella* activity of aqueous and ethanol extract of *Boerhavia diffusa*, tannin and quinone were higher in ethanol extract while saponin was higher in aqueous extract, using agar well diffusion, the aqueous extract showed higher anti-*Salmonella* efficacy while the broth microdilution examined by spectrophotometer revealed that ethanol extract had higher anti-*Salmonella* efficacy. In the present study, the anti-salmonella activity of *Boerhaavia diffusa* may be attributed to an individual or synergistic effect of phytoconstituents present in it. The ethanol and aqueous extracts of leaves of *B. diffusa* whole plant exhibited significant antibacterial activity against both clinical and typed *Salmonella typhi*. Therefore, the plant extract could be used for the treatment of Salmonellosis, however, the in vivo studies is needed to ascertain the safety of the extract.

5. RECOMMENDATION

Based on our findings, it is therefore recommended that both agar well diffusion and broth dilution method should be used to affirm the antimicrobial efficacy of the plant extracts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Prescott LM, Harley JP, Klein DA. Microbiology, McGraw-Hill, Paris. 2005;937-960.

2. Doughari JH. A comparative study on effects of crude extracts of some local medicinal plants and some selected antibiotics on *Salmonella typhi* M. Sc. Thesis Federal University of Technology, Yola; Adamawa State, Nigeria. Yola. 2005; 1-15,1-4.
3. Doughari JH, Pukuma MS, De N. Antibacterial effects of *Balanites aegyptiaca* L. Drel. and *Moringa oleifera* Lam. on *Salmonella typhi*. African Journal of Biotechnology. 2007;6:2212-2215.
4. Birgitta J, Yvonne A, Carl E. Effect of regulation and education and reptile associated salmonellosis. Emerging Infectious Diseases. 2005;11:398-403.
5. Christenson JC. Salmonella infections. Pediatr Rev. 2013;34(9):375-83.
6. Wangari DU. *In vitro* study of anti-Salmonella activities of medicinal plants. Global Journal of Medicinal Plants Research. 2017;3(1):301-308.
7. Kariuki S, Revathi G, Kiiru J, Mengo DM, Mwituria J, Muyodi J, Munyalo A, Yik YT, Kathryn EH, Robert AK, Gordon D. Typhoid in Kenya is associated with dominant multidrug-resistant Salmonella enterica Serovar Typhi haplotype that is also widespread in Southern Asia. Journal of Clinical Microbiology. 2010;48(6):2171-2176.
8. Donald ST, Donatien G, Siméon PCF, Charles F, Fabrice K, Merline ND. *In vivo* anti-salmonella activity of aqueous extract of *Euphorbia prostrata* Aiton (Euphorbiaceae) and its toxicological evaluation. Asian Pacific Journal of Tropical Biomedicine. 2015;5(4):310-318.
9. Bibitha B, Jisha VK, Salitha CV, Mohan S, Valsa AK. Antibacterial activity of different plant extracts. Indian Journal of Microbiology. 2002;42:361-363.
10. Maghrani M, Zeggwagh N, Michel J, Eddouks M. Antihypertensive effect of *Lepidium sativum* L. in spontaneously hypertensive rats. J. Ethnopharm. 2005; 100(102):193-197.

11. Nayak P, Thirunavoukkarasu M. A review of the plant *Boerhaavia diffusa*: Its chemistry, pharmacology and therapeutical potential. The Journal of Phytopharmacology. 2016;5(2):83-92.
12. Katampe I, Uwem M, Uwem J. Antioxidant and antimicrobial effects of aqueous and ethanolic extracts of selected medicinal plants from southern Nigeria. Journal of Advances in Medical and Pharmaceutical Sciences. 2018;17(2):1-12.
13. Girach RD, Singh VK. Notable plants in ethnomedicine of Jajpur District, Orissa, India. In: Recent Progresses in Medicinal Plants. Biopharmaceuticals. 2006;14:495-501.
14. Sourav Das. Antimicrobial activity study of ethanolic extract of *Boerhaavia diffusa* whole plant. International Journal of Pharmacy & Life Sciences. 2012;3(10): 2006-2009.
15. Ogoti P, Magiri E, Magoma G, Kariuki D, Bii C. *In vitro* anti-salmonella activity of extracts from selected Kenyan medicinal plants. Journal of Medicinal Plants Research. 2015;9(8):254-261.
16. Cheesbrough M. Medical laboratory manual for tropical countries. Part II: Microbiology. Cambridge University. USA. 2000;62-70.
17. Marcelin U, Bharathidasan R, Prince L. Antisalmonella activity of selected medicinal plants. World Journal of Pharmacy and Pharmaceutical Sciences. 2016;5(4):2512-2519.
18. Anibijuwon II, Udeze OA. Antimicrobial activity of *Carica papaya* (Pawpaw Leaf) on some pathogenic organisms of clinical origin from South-Western Nigeria. Ethnobotanical Leaflets. 2009;13(7):850-864.
19. Abbas AM, Noor MA, Sarmina Y. Antimicrobial screening of different extracts of *Piper longum* Linn. Res. J Agri and Biol Scien. 2007;3(6):852-857.
20. Shahina N, Samia A, Sheikh AR. *In-vitro* antimicrobial activity of the extracts derived from *Terminalia catappa*. Res. J Microbial. 2007;2(2):180-184.
21. Marjorie MC. Plant products as antimicrobial agents. Clin. Microbiol. Rev. 1999; 12(4):564-582.
22. Abo KA, Ashidi JS. Antimicrobial screening of *Bridelia micrantha*, *Alchornea cordifolia* and *Boerhavia diffusa*. African Journal of Medicine and Medical Sciences. 1999; 28(34):167-169.
23. Madani A, Jain SK. Anti-salmonella activity of *Terminalia bellerica*: *In vitro* and *in vivo* studies. Indian Journal of Experimental Biology. 2008;46:817-821.
24. Olila D, Olwa-Odyek A, Opuda-Asibo J. Antibacterial and antifungal activities of extracts of *Zanthoxylum chabeum* and *Wurbugia ugandensis*, Uganda Medicinal Plants. Afr. Health Sci. 2001;1(2):66-72.
25. Edeoga HO, Okwu DE, Mbaebie O. Nutritional values of some known conventional leafy vegetables of Nigeria. African Journal of Biotechnology. 2005;4: 685-688.
26. Okwute SK. Plant derived pesticidal and antimicrobial agents for use in Agriculture. A review of phytochemical and biological studies on some Nigerian plants. J. Agric. Sci. Technol. 1992;2(1):62-70.
27. Desai SD, Desai DG, Kaur H. Saponins and their biological activities. Pharma Times. 2009;41(3):88-92.
28. Okwu DE. Phytochemical and vitamin content of indigenous species of south-eastern Nigeria. Journal of Sustainable Agriculture and the Environment. 2004;6: 30-37.
29. Shah AS, Wakade AS, Juvekar AR. Immunomodulatory activity of methanolic extract of *Murraya koenigii* Spreng leaves. Indian J Exp Biol. 2008;46:505-9.

© 2019 Oluwabusayo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49728>